

ISSN 2077–9879

Eurasian Mathematical Journal

2013, Volume 4, Number 1

Founded in 2010 by
the L.N. Gumilyov Eurasian National University
in cooperation with
the M.V. Lomonosov Moscow State University
the Peoples' Friendship University of Russia
the University of Padua

Supported by the ISAAC
(International Society for Analysis, its Applications and Computation)

Published by
the L.N. Gumilyov Eurasian National University
Astana, Kazakhstan

EURASIAN MATHEMATICAL JOURNAL

Editorial Board

Editors-in-Chief

V.I. Burenkov, M. Otelbaev, V.A. Sadovnichy

Editors

Sh.A. Alimov (Uzbekistan), H. Begehr (Germany), O.V. Besov (Russia), B. Bójarski (Poland), A.A. Borubaev (Kyrgyzstan), G. Bourdaud (France), R.C. Brown (USA), A. Caetano (Portugal), M. Carro (Spain), A.D.R. Choudary (Pakistan), V.N. Chubarikov (Russia), A.S. Dzumadildaev (Kazakhstan), V.M. Filippov (Russia), H. Ghazaryan (Armenia), M.L. Goldman (Russia), V. Goldshtein (Israel), V. Guliyev (Azerbaijan), D.D. Haroske (Germany), A. Hasanoglu (Turkey), M. Huxley (Great Britain), M. Imanaliev (Kyrgyzstan), P. Jain (India), T.Sh. Kalmenov (Kazakhstan), K.K. Kenzhibaev (Kazakhstan), E. Kissin (Great Britain), V. Kokilashvili (Georgia), A. Kufner (Czech Republic), L.K. Kussainova (Kazakhstan), P.D. Lamberti (Italy), M. Lanza de Cristoforis (Italy), V.G. Maz'ya (Sweden), A.V. Mikhalev (Russia), E.D. Nursultanov (Kazakhstan), R. Oinarov (Kazakhstan), I.N. Parasidis (Greece), J. Pečarić (Croatia), S.A. Plaksa (Ukraine), L.-E. Persson (Sweden), E.L. Presman (Russia), M.D. Ramazanov (Russia), M. Reissig (Germany), S. Sagitov (Sweden), T.O. Shaposhnikova (Sweden), A.A. Shkalikov (Russia), V.A. Skvortsov (Poland), G. Sinnamon (Canada), E.S. Smailov (Kazakhstan), V.D. Stepanov (Russia), Ya.T. Sultanaev (Russia), I.A. Taimanov (Russia), T.V. Tararykova (Great Britain), U.U. Umirbaev (Kazakhstan), Z.D. Usmanov (Tajikistan), B. Viscolani (Italy), Masahiro Yamamoto (Japan), Dachun Yang (China), B.T. Zhumagulov (Kazakhstan)

Managing Editor

A.M. Temirkhanova

Executive Editor

D.T. Matin

Aims and Scope

The Eurasian Mathematical Journal (EMJ) publishes carefully selected original research papers in all areas of mathematics written by mathematicians, principally from Europe and Asia. However papers by mathematicians from other continents are also welcome.

From time to time the EMJ will also publish survey papers.

The EMJ publishes 4 issues in a year.

The language of the paper must be English only.

The contents of EMJ are indexed in Mathematical Reviews, Zentralblatt Math (ZMATH), Referativnyi Zhurnal – Matematika, Math-Net.Ru.

Information for the Authors

Submission. Manuscripts should be written in LaTeX and should be submitted electronically in DVI, PostScript or PDF format to the EMJ Editorial Office via e-mail (eurasianmj@yandex.kz).

When the paper is accepted, the authors will be asked to send the tex-file of the paper to the Editorial Office.

The author who submitted an article for publication will be considered as a corresponding author. Authors may nominate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not guaranteed.

Copyright. When the paper is accepted, the copyright is automatically transferred to the EMJ. Manuscripts are accepted for review on the understanding that the same work has not been already published (except in the form of an abstract), that it is not under consideration for publication elsewhere, and that it has been approved by all authors.

Title page. The title page should start with the title of the paper and authors' names (no degrees). It should contain the Keywords (no more than 10), the Subject Classification (AMS Mathematics Subject Classification (2010) with primary (and secondary) subject classification codes), and the Abstract (no more than 150 words with minimal use of mathematical symbols).

Figures. Figures should be prepared in a digital form which is suitable for direct reproduction.

References. Bibliographical references should be listed alphabetically at the end of the article. The authors should consult the Mathematical Reviews for the standard abbreviations of journals' names.

Authors' data. The authors' affiliations, addresses and e-mail addresses should be placed after the References.

Proofs. The authors will receive proofs only once. The late return of proofs may result in the paper being published in a later issue.

Honorarium. An honorarium will be paid to the authors (at 50 \$ per page, but no more than 2000 \$ for a paper) and to the reviewers (at 50 \$ per review). All papers will be edited with regard to usage of English. If severe editing is required, the honorarium may be reduced to 40 \$ per page. Papers may be submitted in Russian. They will be translated into English, and in this case the honorarium will be 25 \$ per page.

Offprints. The authors will receive offprints in electronic form.

Publication Ethics and Publication Malpractice

Manuscripts are accepted for review on the understanding that the same work has not been already published (except in the form of an abstract), that it is not under consideration for publication elsewhere, and that it has been approved by all authors. Only original papers are accepted for consideration. In particular, translations into English of papers already published in another language are not accepted. No other forms of scientific misconduct are allowed, such as plagiarism, falsification, incorrect interpretation of other works, incorrect citations, etc.

Web-page

The web-page of EMJ is www.emj.enu.kz. One can enter the web-page by typing Eurasian Mathematical Journal in any search engine (Google, Yandex, etc.). The archive of the web-page contains all papers published in EMJ (free access).

Subscription

For Institutions

- US\$ 200 (or equivalent) for one volume (4 issues)
- US\$ 60 (or equivalent) for one issue

For Individuals

- US\$ 160 (or equivalent) for one volume (4 issues)
- US\$ 50 (or equivalent) for one issue.

The price includes handling and postage.

The Subscription Form for subscribers can be obtained by e-mail:

eurasianmj@yandex.kz

The Eurasian Mathematical Journal (EMJ)
The Editorial Office
The L.N. Gumilyov Eurasian National University,
Main Building
Room 355
Tel.: +7-7172-709500 extension 31313
5 Munaitpasov Street
010008 Astana
Kazakhstan

Proceedings of the international workshop
OPERATORS IN MORREY-TYPE SPACES AND APPLICATIONS
(OMTSA 2011)

dedicated to 70th birthday of Professor V.I. Burenkov

held through May 20-27, 2011

at the Ahi Evran University, Kirşehir, Turkey

Volume 2

**WEAK AND STRONG TYPE ESTIMATES
FOR FRACTIONAL INTEGRAL OPERATORS ON MORREY SPACES
OVER METRIC MEASURE SPACES**

I. Sihwaningrum, Y. Sawano

Communicated by T.V. Tararykova

Key words: Weak type estimates, fractional integral operators, Morrey spaces, non-doubling measure.

AMS Mathematics Subject Classification: 42B20, 26A33, 47B38, 47G10.

Abstract. We discuss here a weak and strong type estimate for fractional integral operators on Morrey spaces over metric measure spaces, where the underlying measure does not always satisfy the doubling condition.

1 Introduction

The aim of this paper is to propose a framework of Morrey spaces and fractional integral operators on a metric measure space (X, d, μ) , where μ is a Radon measure.

We recall that the Riesz potential I_α on \mathbb{R}^d is given by

$$I_\alpha f(x) = \int_{\mathbb{R}^d} \frac{f(y)}{|x - y|^{d-\alpha}} dy.$$

According to the Hardy-Littlewood-Sobolev theorem [2, 3, 10], I_α is bounded from $L^p(\mathbb{R}^d)$ to $L^q(\mathbb{R}^d)$ as long as $p, q \in (1, \infty)$ satisfy $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{d}$. Morrey spaces, named after C. Morrey, can also be used to describe the boundedness property of I_α . Here we adopt the following notation to denote Morrey spaces. Let $1 \leq q \leq p < \infty$. For a measurable function f on \mathbb{R}^d , we define

$$\|f\|_{\mathcal{M}_q^p} := \sup \left\{ |B|^{\frac{1}{p} - \frac{1}{q}} \|f\|_{L^q(B)} : B \text{ is a ball} \right\}.$$

The space $\mathcal{M}_q^p(\mathbb{R}^d)$ denotes the set of all measurable functions f for which the norm $\|f\|_{\mathcal{M}_q^p}$ is finite. According to Adams [1], I_α is bounded from $\mathcal{M}_q^p(\mathbb{R}^d)$ to $\mathcal{M}_t^s(\mathbb{R}^d)$, provided that $p, q, s, t \in (1, \infty)$ satisfy $\frac{p}{q} = \frac{t}{s}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{d}$.

In this paper, we aim to show that this theorem is independent from the geometric structure of \mathbb{R}^d by extending it to metric measure spaces, where all we have are the distance function d and the Radon measure μ .

Let (X, d, μ) be a metric measure space with a distance function d and a Borel measure μ . Recall that the measure μ is a doubling measure if it satisfies the so-called *doubling condition*, that is, there exists a constant $C > 0$ such that

$$\mu(B(a, 2r)) \leq C\mu(B(a, r)) \tag{1.1}$$

for every ball $B(a, r)$ with center $a \in X$ and radius $r > 0$. The doubling condition was a key property in classical harmonic analysis but around a decade ago, it turned out to be unnecessary. The point is that we modify the related definitions. Indeed, in the present paper, we propose to redefine the fractional integral operator by

$$I_\alpha f(x) := \int_X \frac{f(y)}{\mu(B(x, 2d(x, y)))^{1-\alpha}} d\mu(y). \quad (1.2)$$

Note that the definition is independent of any notion of dimensions. The same can be said for Morrey spaces, which we define now. For $k > 0, 1 \leq p < \infty$ and $f \in L^1_{\text{loc}}(\mu)$, the norm is given by

$$\begin{aligned} & \|f\|_{\mathcal{M}_1^p(k, \mu)} \\ & := \sup \left\{ \mu(B(x, kr))^{1/p-1} \|\chi_{B(x, r)} f\|_{L^1(\mu)} : x \in X, r > 0, \mu(B(x, r)) > 0 \right\}, \end{aligned}$$

where $\chi_{B(x, r)}$ denotes the characteristic function of the ball $B(x, r)$.

We will prove here that I_α satisfies weak and strong type estimates on Morrey spaces. Our main results are:

Theorem 1.1. *If $1 < p < \infty, 1 < s < \infty, 0 < \alpha < \frac{1}{p}$ and $\frac{1}{s} = \frac{1}{p} - \alpha$, then there exists $C > 0$ such that*

$$\mu\{x \in B(a, r) : I_\alpha f(x) > \gamma\} \leq C \mu(B(a, 6r))^{1-1/p} \left(\frac{\|f\|_{\mathcal{M}_1^p(2, \mu)}}{\gamma} \right)^{s/p}$$

for all positive μ -measurable functions f .

Theorem 1.2. *If $1 < q \leq p < \infty, 1 < s < \infty, 0 < \alpha < \frac{1}{p}, \frac{q}{p} = \frac{t}{s}$ and $\frac{1}{s} = \frac{1}{p} - \alpha$, then there exists $C > 0$ such that*

$$\|I_\alpha f\|_{\mathcal{M}_t^s(6, \mu)} \leq C \|f\|_{\mathcal{M}_q^p(2, \mu)}$$

for all positive μ -measurable functions f .

It hardly looks likely to replace $2d(x, y)$ with $d(x, y)$ in the definition of fractional integral operators and have the similar results according to the example in [9, Section 2]. The proof is a future work.

2 Main Results

We define, for $k > 0$, the centered maximal operator

$$M_k f(x) := \sup_{r>0} \frac{1}{\mu(B(x, kr))} \int_{B(x, r)} |f(y)| d\mu(y) \quad (x \in \text{supp}(\mu)).$$

For the maximal operator M_2 , we prove the following boundedness property on Morrey spaces.

Theorem 2.1. *For any $\gamma > 0$, any positive μ -measurable function k and any ball $B(a, r)$,*

$$\mu\{x \in B(a, r) : M_2 f(x) > \gamma\} \leq 4 \frac{\mu(B(a, 6r))^{1-1/p}}{\gamma} \|f\|_{\mathcal{M}_1^p(2, \mu)}.$$

Proof. We actually prove the estimate

$$\mu\{x \in B(a, r) : M_2 f(x) > 2\gamma\} \leq 2 \frac{\mu(B(a, 6r))^{1-1/p}}{\gamma} \|f\|_{\mathcal{M}_1^p(2, \mu)}. \quad (2.1)$$

Once we prove that

$$\mu\{x \in B(a, r) : M_2[\chi_{B(a, 3r)} f](x) > \gamma\} \leq \frac{\mu(B(a, 6r))^{1-1/p}}{\gamma} \|f\|_{\mathcal{M}_1^p(2, \mu)} \quad (2.2)$$

and

$$\mu\{x \in B(a, r) : M_2[\chi_{X \setminus B(a, 3r)} f](x) > \gamma\} \leq \frac{\mu(B(a, 6r))^{1-1/p}}{\gamma} \|f\|_{\mathcal{M}_1^p(2, \mu)}, \quad (2.3)$$

then estimate (2.1) follows automatically. Estimate (2.2) follows from the weak- $L^1(\mu)$ boundedness of M_2 (see [8, 11]).

Denote by $\mathcal{B}(\mu)$ the set of all balls with positive μ -measure. A geometric observation shows that

$$M_2[\chi_{X \setminus B(a, 3r)} f](x) \leq \sup_{\substack{B \in \mathcal{B}(\mu), B \cap B(a, r) \neq \emptyset, \\ B \cap (X \setminus B(a, 3r)) \neq \emptyset}} \frac{1}{\mu(2B)} \int_B |f(y)| d\mu(y).$$

Let B be a ball which intersects both $B(a, r)$ and $X \setminus B(a, 3r)$. The ball B engulfs $B(a, r)$ if we double the radius of B . Thus,

$$\begin{aligned} & \mu(B(a, 6r))^{1/p-1} \mu\{x \in B(a, r) : M_2[\chi_{B(a, 3r)} f](x) > \gamma\} \\ & \leq \mu(B(a, r))^{1/p-1} \sup_{\substack{B \in \mathcal{B}(\mu), B \cap B(a, r) \neq \emptyset, \\ B \cap (X \setminus B(a, 3r)) \neq \emptyset}} \frac{1}{\mu(2B)} \int_B |f(y)| d\mu(y) \\ & \leq \sup_{\substack{B \in \mathcal{B}(\mu), B \cap B(a, r) \neq \emptyset, \\ B \cap (X \setminus B(a, 3r)) \neq \emptyset}} \frac{\mu(2B)^{1/p}}{\mu(2B)} \int_B |f(y)| d\mu(y) \\ & \leq \|f\|_{\mathcal{M}_1^p(2, \mu)}. \end{aligned}$$

Thus, (2.3) follows. \square

Analogously, the following inequality holds:

Theorem 2.2. *Let $1 < q \leq p < \infty$. Then there exists $C > 0$ such that*

$$\|M_2 f\|_{\mathcal{M}_q^p(6, \mu)} \leq C \|f\|_{\mathcal{M}_q^p(2, \mu)}$$

for all positive μ -measurable functions f .

The proof of Theorem 2.2 being similar to that of Theorem 2.1, we skip the proof, which is based on the $L^q(\mu)$ -boundedness of M_2 established in [8].

Next, we prove a Hedberg type estimate [4].

Theorem 2.3. *If $1 < p < \infty$ and $0 < \alpha < \frac{1}{p}$, then there exists $C > 0$ such that*

$$|I_\alpha f(x)| \leq CM_2 f(x)^{1-p\alpha} \|f\|_{\mathcal{M}_1^p(2,\mu)}^{p\alpha} \quad (x \in X)$$

for all positive μ -measurable functions.

Proof. Let $x \in X$ be fixed. We define

$$R_k(x) := \inf (\{R > 0 : \mu(B(x, 2R)) > 2^k\} \cup \{\infty\}).$$

Then, we have

$$\begin{aligned} & |I_\alpha f(x)| \\ & \leq \sum_{k=-\infty}^{\infty} \lim_{\varepsilon \downarrow 0} \int_{B(x, R_k(x)) \setminus B(x, R_{k-1}(x))} \frac{|f(y)|}{\mu(B(x, 2d(x, y) + \varepsilon))^{1-\alpha}} d\mu(y) \\ & \leq \sum_{k=-\infty}^{\infty} \lim_{\varepsilon \downarrow 0} \int_{B(x, R_k(x)) \setminus B(x, R_{k-1}(x))} \frac{|f(y)|}{\mu(B(x, 2R_{k-1}(x) + \varepsilon))^{1-\alpha}} d\mu(y) \\ & = \sum_{k=-\infty}^{\infty} \lim_{\varepsilon \downarrow 0} \frac{1}{\mu(B(x, 2R_{k-1}(x) + \varepsilon))^{1-\alpha}} \int_{B(x, R_k(x)) \setminus B(x, R_{k-1}(x))} |f(y)| d\mu(y) \\ & \leq \sum_{k \in \mathbb{Z}; R_{k-1}(x) < R_k(x)} \lim_{\varepsilon \downarrow 0} \frac{1}{\mu(B(x, 2R_{k-1}(x) + \varepsilon))^{1-\alpha}} \int_{B(x, R_k(x))} |f(y)| d\mu(y). \end{aligned}$$

The condition $R_{k-1}(x) < R_k(x)$ means that

$$2^{k-1} < \mu(B(x, 2R_{k-1}(x) + \varepsilon)) \leq 2^k$$

for each $\varepsilon \in (0, R_k(x) - R_{k-1}(x))$. Therefore

$$\begin{aligned} |I_\alpha f(x)| & \leq C \sum_{k=-\infty}^{\infty} 2^{k\alpha} \min(M_2 f(x), 2^{-k/p} \|f\|_{\mathcal{M}_1^p(2,\mu)}) \\ & \leq CM f(x)^{1-p\alpha} \|f\|_{\mathcal{M}_1^p(2,\mu)}^{p\alpha}. \end{aligned}$$

Thus, the estimate is proved. □

Now we prove Theorem 1.1.

Proof. For $|I_\alpha f(x)| > \gamma$, Theorem 2.3 gives us

$$M_2 f(x) > \left(\frac{\gamma}{C \|f\|_{\mathcal{M}_1^p(2,\mu)}^{p\alpha}} \right)^{1/(1-p\alpha)}.$$

Hence, by applying Theorem 2.1, we obtain

$$\begin{aligned}
& \mu\{x \in B(a, r) : |I_\alpha f(x)| > \gamma\} \\
& \leq \mu \left\{ x \in B(a, r) : M_2 f(x) > \left(\frac{\gamma}{C \|f\|_{\mathcal{M}_1^p(2, \mu)}^{p\alpha}} \right)^{1/(1-p\alpha)} \right\} \\
& \leq C \mu(B(a, 6r))^{1-1/p} \|f\|_{\mathcal{M}_1^p(2, \mu)} \left(\frac{\|f\|_{\mathcal{M}_1^p(2, \mu)}^{p\alpha}}{\gamma} \right)^{1/(1-p\alpha)} \\
& \leq C \mu(B(a, 6r))^{1-1/p} \frac{\|f\|_{\mathcal{M}_1^p(2, \mu)}^{1+\alpha s}}{\gamma^{s/p}} \\
& \leq C \mu(B(a, 6r))^{1-1/p} \left(\frac{\|f\|_{\mathcal{M}_1^p(2, \mu)}}{\gamma} \right)^{s/p}.
\end{aligned}$$

Thus, the proof is complete. \square

Theorem 1.2 can be proved in a similar way by using Theorem 2.3.

Acknowledgments

The first author was supported by Fundamental Research Program 2012 by Directorate General of Higher Education, Ministry of Education and Culture, Indonesia. The second author was financially supported by Grant-in-Aid for Young Scientists (B), No. 21740104, Japan Society for the Promotion of Science.

This research project is supported by the GCOE program of Kyoto University.

References

- [1] D. Adams, *A note on Riesz potentials*. Duke Math. J. 42 (1975), 765–778.
- [2] G.H. Hardy, J.E. Littlewood, *Some properties of fractional integrals. I*, Math. Zeit. 27 (1927), 565–606.
- [3] G.H. Hardy, J.E. Littlewood, *Some properties of fractional integrals. II*, Math. Zeit. 34 (1932), 403–439.
- [4] L.I. Hedberg, *On certain convolution inequalities*, Proc. Amer. Math. Soc. 36 (1972), 505–510.
- [5] G. Liu, L. Shu, *Boundedness for the commutator of fractional integral on generalized Morrey space in nonhomogenous space*, Anal. Theory Appl. 27 (2011), 51–58.
- [6] E. Nakai, *Hardy-Littlewood maximal operator, singular integral operators, and the Riesz potentials on generalized Morrey spaces*, Math. Nachr. 166 (1994), 95–103.
- [7] F. Nazarov, S. Treil, A. Volberg, *Weak type estimates and Cotlar inequalities for Calderón-Zygmund operators on non-homogeneous spaces*, Internat. Math. Res. Notices, 9 (1998), 463–487.
- [8] Y. Sawano, *Sharp estimates of the modified Hardy-Littlewood maximal operator on the nonhomogeneous space via covering lemmas*, Hokkaido Math. J. 34 (2005), 435–458.
- [9] Y. Sawano, T. Shimomura, *Sobolev embeddings for Riesz potentials of functions in non-doubling Morrey spaces of variable exponents*. Collectanea Mathematica, online.
- [10] S.L. Sobolev, *On a theorem in functional analysis*. (in Russian), Mat. Sb. 46 (1938), 471–497 [English translation in Amer. Math. Soc. Transl. ser. 2, 34 (1963), 39–68.
- [11] Y. Terasawa, *Outer measures and weak type (1,1) estimates of Hardy-Littlewood maximal operators*, J. Inequal. Appl., 2006, Art. ID 15063, 13 pp.

Idha Sihwaningrum
Faculty of Sciences and Engineering
Jenderal Soedirman University
Purwokerto, 53122 Indonesia
E-mail: idha.sihwaningrum@unsoed.ac.id

Yoshihiro Sawano
Department of Mathematics and Information Sciences
Tokyo Metropolitan University
Tokyo 192-0397, Japan
E-mail: ysawano@tmu.ac.jp

Received: 15.11.2012