

ISSN (Print): 2077-9879
ISSN (Online): 2617-2658

Eurasian Mathematical Journal

2020, Volume 11, Number 4

Founded in 2010 by
the L.N. Gumilyov Eurasian National University
in cooperation with
the M.V. Lomonosov Moscow State University
the Peoples' Friendship University of Russia (RUDN University)
the University of Padua

Starting with 2018 co-funded
by the L.N. Gumilyov Eurasian National University
and
the Peoples' Friendship University of Russia (RUDN University)

Supported by the ISAAC
(International Society for Analysis, its Applications and Computation)
and
by the Kazakhstan Mathematical Society

Published by
the L.N. Gumilyov Eurasian National University
Nur-Sultan, Kazakhstan

EURASIAN MATHEMATICAL JOURNAL

Editorial Board

Editors-in-Chief

V.I. Burenkov, M. Otelbaev, V.A. Sadovnichy

Vice-Editors-in-Chief

K.N. Ospanov, T.V. Tararykova

Editors

Sh.A. Alimov (Uzbekistan), H. Begehr (Germany), T. Bekjan (China), O.V. Besov (Russia), N.K. Blied (Kazakhstan), N.A. Bokayev (Kazakhstan), A.A. Borubaev (Kyrgyzstan), G. Bourdaud (France), A. Caetano (Portugal), M. Carro (Spain), A.D.R. Choudary (Pakistan), V.N. Chubarikov (Russia), A.S. Dzumadildaev (Kazakhstan), V.M. Filippov (Russia), H. Ghazaryan (Armenia), M.L. Goldman (Russia), V. Goldshtein (Israel), V. Guliyev (Azerbaijan), D.D. Haroske (Germany), A. Hasanoglu (Turkey), M. Huxley (Great Britain), P. Jain (India), T.Sh. Kalmenov (Kazakhstan), B.E. Kangyzhin (Kazakhstan), K.K. Kenzhibayev (Kazakhstan), S.N. Kharin (Kazakhstan), E. Kissin (Great Britain), V. Kokilashvili (Georgia), V.I. Korzyuk (Belarus), A. Kufner (Czech Republic), L.K. Kussainova (Kazakhstan), P.D. Lamberti (Italy), M. Lanza de Cristoforis (Italy), F. Lanzara (Italy), V.G. Maz'ya (Sweden), K.T. Mynbayev (Kazakhstan), E.D. Nursultanov (Kazakhstan), R. Oinarov (Kazakhstan), I.N. Parasidis (Greece), J. Pečarić (Croatia), S.A. Plaksa (Ukraine), L.-E. Persson (Sweden), E.L. Presman (Russia), M.A. Ragusa (Italy), M.D. Ramazanov (Russia), M. Reissig (Germany), M. Ruzhansky (Great Britain), M.A. Sadybekov (Kazakhstan), S. Sagitov (Sweden), T.O. Shaposhnikova (Sweden), A.A. Shkalikov (Russia), V.A. Skvortsov (Poland), G. Sinamon (Canada), E.S. Smailov (Kazakhstan), V.D. Stepanov (Russia), Ya.T. Sultanaev (Russia), D. Suragan (Kazakhstan), I.A. Taimanov (Russia), J.A. Tussupov (Kazakhstan), U.U. Umirbaev (Kazakhstan), Z.D. Usmanov (Tajikistan), N. Vasilevski (Mexico), Dachun Yang (China), B.T. Zhumagulov (Kazakhstan)

Managing Editor

A.M. Temirkhanova

Aims and Scope

The Eurasian Mathematical Journal (EMJ) publishes carefully selected original research papers in all areas of mathematics written by mathematicians, principally from Europe and Asia. However papers by mathematicians from other continents are also welcome.

From time to time the EMJ publishes survey papers.

The EMJ publishes 4 issues in a year.

The language of the paper must be English only.

The contents of the EMJ are indexed in Scopus, Web of Science (ESCI), Mathematical Reviews, MathSciNet, Zentralblatt Math (ZMATH), Referativnyi Zhurnal – Matematika, Math-Net.Ru.

The EMJ is included in the list of journals recommended by the Committee for Control of Education and Science (Ministry of Education and Science of the Republic of Kazakhstan) and in the list of journals recommended by the Higher Attestation Commission (Ministry of Education and Science of the Russian Federation).

Information for the Authors

Submission. Manuscripts should be written in LaTeX and should be submitted electronically in DVI, PostScript or PDF format to the EMJ Editorial Office through the provided web interface (www.enu.kz).

When the paper is accepted, the authors will be asked to send the tex-file of the paper to the Editorial Office.

The author who submitted an article for publication will be considered as a corresponding author. Authors may nominate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not guaranteed.

Copyright. When the paper is accepted, the copyright is automatically transferred to the EMJ. Manuscripts are accepted for review on the understanding that the same work has not been already published (except in the form of an abstract), that it is not under consideration for publication elsewhere, and that it has been approved by all authors.

Title page. The title page should start with the title of the paper and authors' names (no degrees). It should contain the Keywords (no more than 10), the Subject Classification (AMS Mathematics Subject Classification (2010) with primary (and secondary) subject classification codes), and the Abstract (no more than 150 words with minimal use of mathematical symbols).

Figures. Figures should be prepared in a digital form which is suitable for direct reproduction.

References. Bibliographical references should be listed alphabetically at the end of the article. The authors should consult the Mathematical Reviews for the standard abbreviations of journals' names.

Authors' data. The authors' affiliations, addresses and e-mail addresses should be placed after the References.

Proofs. The authors will receive proofs only once. The late return of proofs may result in the paper being published in a later issue.

Offprints. The authors will receive offprints in electronic form.

Publication Ethics and Publication Malpractice

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the EMJ implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The EMJ follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (<http://publicationethics.org/files/u2/NewCode.pdf>). To verify originality, your article may be checked by the originality detection service CrossCheck <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the EMJ.

The Editorial Board of the EMJ will monitor and safeguard publishing ethics.

The procedure of reviewing a manuscript, established by the Editorial Board of the Eurasian Mathematical Journal

1. Reviewing procedure

1.1. All research papers received by the Eurasian Mathematical Journal (EMJ) are subject to mandatory reviewing.

1.2. The Managing Editor of the journal determines whether a paper fits to the scope of the EMJ and satisfies the rules of writing papers for the EMJ, and directs it for a preliminary review to one of the Editors-in-chief who checks the scientific content of the manuscript and assigns a specialist for reviewing the manuscript.

1.3. Reviewers of manuscripts are selected from highly qualified scientists and specialists of the L.N. Gumilyov Eurasian National University (doctors of sciences, professors), other universities of the Republic of Kazakhstan and foreign countries. An author of a paper cannot be its reviewer.

1.4. Duration of reviewing in each case is determined by the Managing Editor aiming at creating conditions for the most rapid publication of the paper.

1.5. Reviewing is confidential. Information about a reviewer is anonymous to the authors and is available only for the Editorial Board and the Control Committee in the Field of Education and Science of the Ministry of Education and Science of the Republic of Kazakhstan (CCFES). The author has the right to read the text of the review.

1.6. If required, the review is sent to the author by e-mail.

1.7. A positive review is not a sufficient basis for publication of the paper.

1.8. If a reviewer overall approves the paper, but has observations, the review is confidentially sent to the author. A revised version of the paper in which the comments of the reviewer are taken into account is sent to the same reviewer for additional reviewing.

1.9. In the case of a negative review the text of the review is confidentially sent to the author.

1.10. If the author sends a well reasoned response to the comments of the reviewer, the paper should be considered by a commission, consisting of three members of the Editorial Board.

1.11. The final decision on publication of the paper is made by the Editorial Board and is recorded in the minutes of the meeting of the Editorial Board.

1.12. After the paper is accepted for publication by the Editorial Board the Managing Editor informs the author about this and about the date of publication.

1.13. Originals reviews are stored in the Editorial Office for three years from the date of publication and are provided on request of the CCFES.

1.14. No fee for reviewing papers will be charged.

2. Requirements for the content of a review

2.1. In the title of a review there should be indicated the author(s) and the title of a paper.

2.2. A review should include a qualified analysis of the material of a paper, objective assessment and reasoned recommendations.

2.3. A review should cover the following topics:

- compliance of the paper with the scope of the EMJ;
- compliance of the title of the paper to its content;
- compliance of the paper to the rules of writing papers for the EMJ (abstract, key words and phrases, bibliography etc.);
- a general description and assessment of the content of the paper (subject, focus, actuality of the topic, importance and actuality of the obtained results, possible applications);
- content of the paper (the originality of the material, survey of previously published studies on the topic of the paper, erroneous statements (if any), controversial issues (if any), and so on);

- exposition of the paper (clarity, conciseness, completeness of proofs, completeness of bibliographic references, typographical quality of the text);
- possibility of reducing the volume of the paper, without harming the content and understanding of the presented scientific results;
- description of positive aspects of the paper, as well as of drawbacks, recommendations for corrections and complements to the text.

2.4. The final part of the review should contain an overall opinion of a reviewer on the paper and a clear recommendation on whether the paper can be published in the Eurasian Mathematical Journal, should be sent back to the author for revision or cannot be published.

Web-page

The web-page of the EMJ is www.emj.enu.kz. One can enter the web-page by typing Eurasian Mathematical Journal in any search engine (Google, Yandex, etc.). The archive of the web-page contains all papers published in the EMJ (free access).

Subscription

Subscription index of the EMJ 76090 via KAZPOST.

E-mail

eurasianmj@yandex.kz

The Eurasian Mathematical Journal (EMJ)
The Nur-Sultan Editorial Office
The L.N. Gumilyov Eurasian National University
Building no. 3
Room 306a
Tel.: +7-7172-709500 extension 33312
13 Kazhymukan St
010008 Nur-Sultan, Kazakhstan

The Moscow Editorial Office
The Peoples' Friendship University of Russia
(RUDN University)
Room 562
Tel.: +7-495-9550968
3 Ordzonikidze St
117198 Moscow, Russia

UNCONDITIONAL BASES OF SYSTEMS OF BESSEL FUNCTIONS

B.V. Vynnyts'kyi, R.V. Khats', I.B. Sheparovych

Communicated by E.S. Smailov

Key words: interpolation problem, complete interpolating sequence, unconditional basis, Bessel function, entire function of exponential type.

AMS Mathematics Subject Classification: Primary 41A05, 30E05, 30B60, 33C10, 34B30, 41A30, 42A65; Secondary 30D10, 30D20, 44A15, 46E30.

Abstract. We find a criterion of unconditional basicity of the system $(\sqrt{x\rho_k}J_\nu(x\rho_k) : k \in \mathbb{N})$ in the space $L^2(0; 1)$ where J_ν is the Bessel function of the first kind of index $\nu \geq -1/2$ and $(\rho_k : k \in \mathbb{N})$ is a sequence of distinct nonzero complex numbers.

DOI: <https://doi.org/10.32523/2077-9879-2020-11-4-76-86>

1 Introduction and main results

Let $P = \{\rho_k : k \in \mathbb{N}\}$ be a sequence of nonzero complex numbers. Developing the results of Pavlov [21], Nikol'skii [20] and others (see [14, 16, 24]), Minkin [19] obtained a criterion of unconditional basicity of the system of exponentials $(\exp(i\rho_k t) : k \in \mathbb{N})$ in $L^2(-\pi; \pi)$. Lyubarskii and Seip [17] found another approach to the proof of this criterion. Let

$$J_\nu(z) = \sum_{k=0}^{\infty} \frac{(-1)^k (z/2)^{\nu+2k}}{k! \Gamma(\nu + k + 1)}, \quad z = x + iy = re^{i\theta},$$

be the Bessel function of the first kind of index $\nu \in \mathbb{R}$, where Γ is the gamma function. In this paper, we will establish a criterion of unconditional basicity of the system

$$(\sqrt{x\rho_k}J_\nu(x\rho_k) : k \in \mathbb{N}) \tag{1.1}$$

in the space $L^2(0; 1)$ if $\nu \geq -1/2$.

Basis properties of the systems of Bessel functions and similar systems have been studied in a number of papers (see, for instance, [2], [12], [13] and [26]–[33]). In particular, it is well known that if $\nu > -1$ and $(\rho_k : k \in \mathbb{N})$ is a sequence of positive zeros of J_ν , then [13] (see also [26], [27], [33]) system (1.1) forms a basis in $L^2(0; 1)$. Another sufficient conditions of basicity of system (1.1) with $\nu \geq -1/2$ in $L^2(0; 1)$ were found in [30].

Let $\lambda_k = \rho_k$, $\lambda_{-k} := -\lambda_k$, $\Lambda = \{\lambda_k : k \in \mathbb{Z} \setminus \{0\}\}$, let $L^2(X)$ be the space of all measurable functions $f : X \rightarrow \mathbb{C}$, $X \subseteq \mathbb{R}$, satisfying

$$\|f\|_{L^2(X)}^2 := \int_X |f(x)|^2 dx < +\infty,$$

let $\nu \in (-1; +\infty)$, let $L^{2,\nu}(\mathbb{R})$ be the space of all Lebesgue measurable functions f satisfying

$$\|f\|_{L^{2,\nu}(\mathbb{R})}^2 := \int_{-\infty}^{+\infty} |x|^{2\nu+1} |f(x)|^2 dx < +\infty,$$

let $PW^{2,\nu}$ be the space of all entire functions f of exponential type $\sigma \leq 1$ for which

$$\|f\|_{PW^{2,\nu}}^2 := \int_{-\infty}^{+\infty} |x|^{2\nu+1} |f(x)|^2 dx < +\infty,$$

and let $PW_+^{2,\nu}$ be the subspace of even functions $f \in PW^{2,\nu}$.

Our main result is the following statement.

Theorem 1.1. *Let $\nu \geq -1/2$. System (1.1) forms an unconditionally basis in $L^2(0;1)$ if and only if the following conditions hold:*

- 1) $\rho_k^2 \neq \rho_j^2$ for $k \neq j$;
- 2) the function

$$S(z) = \prod_{k=1}^{\infty} \left(1 - \frac{z^2}{\rho_k^2}\right)$$

is an entire function of exponential type $\sigma \leq 1$ and $(z^2 - \rho_1^2)^{-1}S(z) \in PW_+^{2,\nu}$;

$$3) \inf \left\{ \prod_{\substack{\text{Im } \lambda_j > 0 \\ j \neq k}} \left| \frac{\lambda_k - \lambda_j}{\lambda_k - \bar{\lambda}_j} \right| : \text{Im } \lambda_k > 0 \right\} > 0, \quad \inf \left\{ \prod_{\substack{\text{Im } \lambda_j < 0 \\ j \neq k}} \left| \frac{\lambda_k - \lambda_j}{\lambda_k - \bar{\lambda}_j} \right| : \text{Im } \lambda_k < 0 \right\} > 0;$$

$$4) \inf \left\{ \frac{|\lambda_k - \lambda_j|}{1 + |\lambda_k - \bar{\lambda}_j|} : (k; j) \in (\mathbb{Z} \setminus \{0\})^2, k \neq j \right\} > 0;$$

5) the function $u(x) = F^2(x)$, $F(x) := |x|^{\nu+1/2} \frac{|S(x)|}{\text{dist}(x; \Lambda)}$, satisfies the (continuous) (A_2) condition

$$\sup_I \left\{ \left(\frac{1}{|I|} \int_I u(x) dx \right) \left(\frac{1}{|I|} \int_I u^{-1}(x) dx \right) : I \subset \mathbb{R} \right\} < +\infty,$$

where $I \subset \mathbb{R}$ is an interval of length $|I|$, and $\text{dist}(x; \Lambda)$ is the distance between the element x and the set Λ .

Remark 1. The conditions 3) – 5) may be expressed in different ways (see [17]). We will discuss this in detail below.

Let $w = (w_k)$ be a sequence of positive numbers, let $l^{2,w}$ be the space of all sequences $d = (d_k)$ for which $\|d\|_{l^{2,w}}^2 := \sum_k |d_k|^2 w_k < +\infty$, and let $l_+^{2,w}$ be the subspace of $l^{2,w}$ of sequences $d = (d_k : k \in \mathbb{Z} \setminus \{0\})$ such that $d_{-k} = d_k$. Interpolation problems in the spaces $PW^{2,\nu}$ have been considered in [3]–[6], [9]–[11], [16], [18] and [22]–[24]. Following Lyubarskii and Seip [17], we say that a sequence $\Lambda = \{\lambda_k : k \in \mathbb{Z} \setminus \{0\}\}$ (sequence $P = \{\rho_k : k \in \mathbb{N}\}$) is a *complete interpolating sequence* for $PW^{2,\nu}$ (for $PW_+^{2,\nu}$) if for every sequence $d \in l^{2,w}$, $w_k := |\lambda_k|^{2\nu+1}(1 + |\text{Im } \lambda_k|)e^{-2|\text{Im } \lambda_k|}$ (sequence $d \in l_+^{2,w}$, $w_k := |\rho_k|^{2\nu+1}(1 + |\text{Im } \rho_k|)e^{-2|\text{Im } \rho_k|}$), the interpolation problem $f(\lambda_k) = d_k$, $k \in \mathbb{Z} \setminus \{0\}$ (interpolation problem $f(\rho_k) = d_k$, $k \in \mathbb{N}$) has a unique solution $f \in PW^{2,\nu}$ ($f \in PW_+^{2,\nu}$). Following [17], we also say that a sequence $w = (w_k)$ satisfies the *discrete (A_2) condition* if

$$\sup \left\{ \left(\frac{1}{n} \sum_{j=k+1}^{k+n} w_j \right) \left(\frac{1}{n} \sum_{j=k+1}^{k+n} w_j^{-1} \right) : n \in \mathbb{N}, k \in \mathbb{Z} \setminus \{0\} \right\} < +\infty.$$

Theorem 1.1 will be proved by using the results of Lyubarskii and Seip ([17, 18]), and the following statement.

Theorem 1.2. *Let $\nu \geq -1/2$ and $w_k = |\rho_k|^{2\nu+1}(1 + |\text{Im } \rho_k|)e^{-2|\text{Im } \rho_k|}$. System (1.1) forms an unconditionally basis in $L^2(0;1)$ if and only if the sequence $P = \{\rho_k : k \in \mathbb{N}\}$ is a complete interpolating sequence for $PW_+^{2,\nu}$.*

2 Proof of Theorem 1.2

To prove Theorem 1.2 we need the following auxiliary statements.

Lemma 2.1. (see [3]) *Let $\nu > -1$. Then every function $f \in L^2(0; +\infty)$ can be represented in the form*

$$f(z) = \int_0^{+\infty} \sqrt{zt} J_\nu(zt) h_f(t) dt$$

with some function $h_f \in L^2(0; +\infty)$. Also, we have $\|f\|_{L^2(0; +\infty)} = \|h_f\|_{L^2(0; +\infty)}$ and

$$h_f(t) = \int_0^{+\infty} \sqrt{zt} J_\nu(zt) f(z) dz.$$

Lemma 2.2. (see [1], [8]) *Let $\nu \geq -1/2$. A function f has the representation*

$$f(z) = \int_0^1 \sqrt{tz} J_\nu(zt) \gamma_f(t) dt$$

with some function $\gamma_f \in L^2(0; 1)$ if and only if $f \in L^2(0; +\infty)$ and $f(z) = z^{\nu+1/2} Q_f(z)$, where Q_f is an even entire function of exponential type $\sigma \leq 1$.

By c_j we denote some positive constants.

Lemma 2.3. *Let $\nu > -1$ and $(\rho_k : k \in \mathbb{N})$ be an arbitrary sequence of nonzero complex numbers. Then there exists a positive constant c_1 such that for all $k \in \mathbb{N}$*

$$e^{2|\operatorname{Im} \rho_k|} (1 + |\operatorname{Im} \rho_k|)^{-1} / c_1 \leq \|\sqrt{t \rho_k} J_\nu(t \rho_k)\|_{L^2(0; 1)}^2 \leq c_1 e^{2|\operatorname{Im} \rho_k|} (1 + |\operatorname{Im} \rho_k|)^{-1}.$$

Proof. In fact, the right-hand side of this inequality follows from the following estimate (see [13], [29], [33])

$$|\sqrt{z} J_\nu(z)| \leq c_2 e^{|\operatorname{Im} z|} \left(\frac{|z|}{1 + |z|} \right)^{\nu+1/2}, \quad z \in \mathbb{C}.$$

Let us prove the left-hand side of this inequality (in the case $\operatorname{Im} \rho_k = 0$ the proof is given in [26, p. 227]). Without loss of generality, we may assume that $|\rho_k| > 1$. Using relations (see [26], [27], [33])

$$J_\nu(z) = \frac{z^\nu}{2^\nu \Gamma(\nu + 1)} + O(z^{\nu+2}), \quad z \rightarrow 0,$$

$$J_\nu(z) = \sqrt{\frac{2}{\pi z}} \cos\left(z - \frac{\pi}{2}\nu - \frac{\pi}{4}\right) + O\left(\frac{e^{|\operatorname{Im} z|}}{|z|^{3/2}}\right), \quad z \rightarrow \infty, \quad |\arg z| < \pi,$$

we get

$$\begin{aligned} \|\sqrt{t \rho_k} J_\nu(t \rho_k)\|_{L^2(0; 1)} &= \left(\int_0^1 |\rho_k| t |J_\nu(t |\rho_k| e^{i\theta_k})|^2 dt \right)^{1/2} = \left(\frac{1}{|\rho_k|} \int_0^{|\rho_k|} t |J_\nu(t e^{i\theta_k})|^2 dt \right)^{1/2} \\ &= \left(\frac{1}{|\rho_k|} \int_0^1 t |J_\nu(t e^{i\theta_k})|^2 dt + \frac{1}{|\rho_k|} \int_1^{|\rho_k|} t |J_\nu(t e^{i\theta_k})|^2 dt \right)^{1/2} \\ &\geq \frac{1}{\sqrt{|\rho_k|}} \left(\int_1^{|\rho_k|} t |J_\nu(t e^{i\theta_k})|^2 dt \right)^{1/2} = \frac{1}{\sqrt{|\rho_k|}} \|\sqrt{t} J_\nu(t e^{i\theta_k})\|_{L^2(1; |\rho_k|)} \\ &\geq \sqrt{\frac{2}{\pi}} \frac{1}{\sqrt{|\rho_k|}} \left\| \cos\left(t e^{i\theta_k} - \frac{\pi}{2}\nu - \frac{\pi}{4}\right) \right\|_{L^2(1; |\rho_k|)} \\ &\quad - \frac{1}{\sqrt{|\rho_k|}} \left\| \frac{e^{t|\sin \theta_k|}}{t} \right\|_{L^2(1; |\rho_k|)}, \quad \rho_k := |\rho_k| e^{i\theta_k}. \end{aligned}$$

In addition, (here and in what follows the sign \asymp means that the ratio of the two sides lies between two positive constants)

$$\begin{aligned} \left\| \frac{e^{t|\sin \theta_k|}}{t} \right\|_{L^2(1;|\rho_k|)}^2 &\asymp \frac{e^{2|\rho_k|\sin \theta_k}}{2|\rho_k|(1+|\rho_k|\sin \theta_k)}, \quad k \rightarrow \infty, \\ |\cos(x+iy)|^2 &= \cos^2 x + \sinh^2 y, \\ \int_1^{|\rho_k|} \sinh^2(t \sin \theta_k) dt &= \frac{\sinh(2|\rho_k|\sin \theta_k) - \sinh(2 \sin \theta_k)}{4 \sin \theta_k} - \frac{|\rho_k| - 1}{2}, \\ \int_1^{|\rho_k|} \cos^2 \left(t \cos \theta_k - \frac{\pi}{2} \nu - \frac{\pi}{4} \right) dt \\ &= \frac{|\rho_k| - 1}{2} + \frac{\sin(|\rho_k| \cos \theta_k - \cos \theta_k) \sin(|\rho_k| \cos \theta_k + \cos \theta_k - \pi \nu)}{2 \cos \theta_k}. \end{aligned}$$

□

Proof of Theorem 1.2. The sequence (e_k) of nonzero elements of a separable Hilbert space H with the inner product $\langle \cdot; \cdot \rangle$ is an unconditional basis of H (see, for example, [25]) if and only if for each sequence (b_k) satisfying $\sum_k |b_k|^2 \|e_k\|^{-2} < +\infty$ there is a unique element $g \in H$ such that $\langle g; e_k \rangle = b_k$ for all $k \in \mathbb{N}$. If $H = L^2(0; 1)$, $e_k = \sqrt{\rho_k} x J_\nu(\rho_k x)$ and $g \in L^2(0; 1)$, then by Lemmas 2.1 and 2.2, we have

$$\langle g; e_k \rangle = \int_0^1 \sqrt{\rho_k} t J_\nu(\rho_k t) \overline{g(t)} dt = \rho_k^{\nu+1/2} Q_f(\rho_k),$$

where $Q_f \in PW_+^{2,\nu}$. Therefore, using Lemma 2.3, we obtain the required statement. □

3 Proof of Theorem 1.1

Theorem 1.1 follows from Theorem 1.2 and the results of Lyubarskii and Seip ([17], [18]). So, we are going to sketch the proof of Theorem 1.1.

Let $-\infty \leq a < b \leq +\infty$ and $\mathbb{C}_{a,b} = \{z : a < \text{Im } z < b\}$, and let $H^2(\mathbb{C}_{a,b})$ be the space of all functions f that are holomorphic in the strip $\mathbb{C}_{a,b}$ and satisfy

$$\sup \left\{ \int_{-\infty}^{+\infty} |f(x+iy)|^2 dx : a < y < b \right\} < +\infty.$$

If at least one of the numbers a or b is finite, then every function $f \in H^2(\mathbb{C}_{a,b})$ has angular limit values $f \in L^2(\partial\mathbb{C}_{a,b})$ almost everywhere on $\partial\mathbb{C}_{a,b}$, and the equality $\|f\|^2 := \int_{\partial\mathbb{C}_{a,b}} |f(z)|^2 |dz|$ determines a norm on $H^2(\mathbb{C}_{a,b})$ (see [3], [15]). If $a \in \mathbb{R}$ and $b = +\infty$, then $H^2(\mathbb{C}_{a,b})$ is the Hardy space in the half-plane $\mathbb{C}_{a,b}$. The same can be said when $a = -\infty$ and $b \in \mathbb{R}$.

Lemma 3.1. (see [3]–[6], [22], [23]) *If $\nu \geq -1/2$ and $f \in PW^{2,\nu}$, then for any $a \in \mathbb{R}$ the function $f_+(z) = (z+ia)^{\nu+1/2} e^{iz} f(z)$ belongs to $H^2(\mathbb{C}_{-a,+\infty})$ and*

$$\int_{-\infty}^{+\infty} |x+iy+ia|^{2\nu+1} |f(x+iy+ia)|^2 dx \leq e^{2y} \int_{-\infty}^{+\infty} |x+ia|^{2\nu+1} |f(x+ia)|^2 dx, \quad y > -a.$$

Moreover, the function $f_-(z) = (z-ia)^{\nu+1/2} e^{-iz} f(z)$ belongs to $H^2(\mathbb{C}_{-\infty,a})$ and

$$\int_{-\infty}^{+\infty} |x+iy-ia|^{2\nu+1} |f(x+iy-ia)|^2 dx \leq e^{-2y} \int_{-\infty}^{+\infty} |x-ia|^{2\nu+1} |f(x-ia)|^2 dx, \quad y < a.$$

Lemma 3.2. (see [3]–[6], [22], [23]) *Let $\nu \geq -1/2$, $\gamma \in \mathbb{R}$ and $\beta \in \mathbb{R}$. The space (as a set) $PW_+^{2,\nu}$ coincides with the set $W^{2,\nu}[\gamma;\beta]$ of all even entire functions f of exponential type $\sigma \leq 1$ satisfying*

$$\|f\|_{W^{2,\nu}[\gamma;\beta]}^2 := \int_{-\infty}^{+\infty} |x + i\gamma|^{2\nu+1} |f(x + i\beta)|^2 dx < +\infty.$$

Moreover, the norms $\|f\|_{W^{2,\nu}[\gamma;\beta]}$ and $\|f\|_{PW_+^{2,\nu}}$ are equivalent,

$$\int_{-\infty}^{+\infty} |x|^{2\nu+1} |f(x + iy)|^2 dx \leq c_3 \|f\|_{PW_+^{2,\nu}} e^{2|y|},$$

and for any $z = x + iy \in \mathbb{C}$ holds

$$|f(z)| \leq c_4 \|f\|_{PW_+^{2,\nu}} e^{|y|} (1 + |z|)^{-\nu-1/2} (1 + |y|)^{-1/2}.$$

Lemma 3.3. (see [17, pp. 362, 363, 366, 367]) *Let $\nu \geq -1/2$. If the sequence $P = \{\rho_k : k \in \mathbb{N}\}$ is a complete interpolating sequence for $PW_+^{2,\nu}$, then for every $k \in \mathbb{Z} \setminus \{0\}$ the interpolation problems $f_k(\lambda_k) = f_k(\lambda_{-k}) = 1$ and $f_k(\lambda_j) = f_k(\lambda_{-j}) = 0$, $j \in \mathbb{Z} \setminus \{0; k; -k\}$, are solvable in $H^2(\mathbb{C}_{0,+\infty})$ and in $H^2(\mathbb{C}_{-\infty,0})$. Moreover, properties 1), 2) and the following ones hold:*

$$\inf \left\{ \prod_{\substack{\text{Im } \lambda_j > a, \\ j \neq k}} \left| \frac{\lambda_k - \lambda_j}{\lambda_k - \bar{\lambda}_j + 2ia} \right| : \text{Im } \lambda_k > a \right\} > 0 \quad \text{for each } a \in \mathbb{R}; \quad (3.1)$$

$$\inf \left\{ \prod_{\substack{\text{Im } \lambda_j < a, \\ j \neq k}} \left| \frac{\lambda_k - \lambda_j}{\lambda_k - \bar{\lambda}_j - 2ia} \right| : \text{Im } \lambda_k < a \right\} > 0 \quad \text{for each } a \in \mathbb{R}; \quad (3.2)$$

$$\sup \left\{ \sum_{\substack{j \in \mathbb{Z} \setminus \{0\}, \\ j \neq k}} \frac{(1 + |\text{Im } \lambda_k|)(1 + |\text{Im } \lambda_j|)}{|\lambda_k - \bar{\lambda}_j|^2} : k \in \mathbb{Z} \setminus \{0\} \right\} < +\infty; \quad (3.3)$$

for some $\varepsilon > 0$ the disks

$$K(\lambda_k) := \{z : |z - \lambda_k| \leq 10\varepsilon(1 + |\text{Im } \lambda_k|)\} \quad \text{are pairwise disjoint}; \quad (3.4)$$

$$\mu_\Lambda := \sum_{\text{Im } \lambda_k \geq 0} \text{Im } \lambda_k \delta_{\lambda_k} \quad (3.5)$$

is a Carleson measure in $\mathbb{C}^+ := \mathbb{C}_{0,+\infty}$ (δ_λ is the unit point measure at λ).

By manipulating the Carleson conditions (3.1) and (3.2) in much the same way as in [7, pp. 288–290], we obtain the following lemma.

Lemma 3.4. (see [17, pp. 363, 364, 367]) *Condition (3.3) is equivalent to conditions 3) and 4), and also to conditions (3.1) and (3.2).*

Lemma 3.5. *Let $\nu \geq -1/2$. If a sequence Λ satisfies condition (3.3), then*

$$\sum_{k \in \mathbb{Z} \setminus \{0\}} (1 + |\text{Im } \lambda_k|)(1 + |\lambda_k|)^{2\nu+1} e^{-2|\text{Im } \lambda_k|} |f(\lambda_k)|^2 \leq c_5 \|f\|_{PW_+^{2,\nu}}^2, \quad f \in PW_+^{2,\nu}.$$

Proof. Indeed, if $f \in PW_+^{2,\nu}$, then by Lemma 3.1, we have $f_+(z) = (z+2i)^{\nu+1/2}e^{iz}f(z) \in H^2(\mathbb{C}_{-2,+\infty})$ and $f_-(z) = (z-2i)^{\nu+1/2}e^{-iz}f(z) \in H^2(\mathbb{C}_{-\infty,2})$, and for the Hardy spaces the corresponding result is true (see [15]). \square

Let $Q(x; r)$ be the square with center at $x \in \mathbb{R}$, side length $2r$, and sides parallel to the coordinate axes. According to [17], we say that a set $P \subset \mathbb{C}$ is *relatively dense* if there exists $r_0 > 0$ such that $P \cap Q(x; r_0) \neq \emptyset$ for each $x \in \mathbb{R}$.

Lemma 3.6. *Let $\nu \geq -1/2$. If $P = \{\rho_k : k \in \mathbb{N}\}$ is a complete interpolating sequence for $PW_+^{2,\nu}$, then $\Lambda = \{\lambda_k : k \in \mathbb{Z} \setminus \{0\}\}$ is relatively dense and for any function $f \in PW_+^{2,\nu}$*

$$\|f\|_{PW_+^{2,\nu}}^2/c_6 \leq \sum_{k \in \mathbb{Z} \setminus \{0\}} (1 + |\operatorname{Im} \lambda_k|)(1 + |\lambda_k|)^{2\nu+1} e^{-2|\operatorname{Im} \lambda_k|} |f(\lambda_k)|^2 \leq c_6 \|f\|_{PW_+^{2,\nu}}^2. \quad (3.6)$$

Proof. The proof of condition (3.6) is the same as that of [17]. Assume that a set Λ is not relatively dense. Then there exist sequences $\{x_j\}$ and $\{r_j\}$ such that $|x_j| > s$, $s \in \mathbb{N}$, $r_j \rightarrow +\infty$ and $\Lambda \cap Q(x_j; r_j) = \emptyset$. Let $s > 1 + \nu$, $\mu = \nu + 1 - s$, $q_j = 1/|x_j|$ and let $f_j(z) = q_j^\mu \left(\sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} q_j^{2k+1} (z^2 - x_j^2)^k \right)^s$. Then $f_j \in PW_+^{2,\nu}$,

$$f_j(z) = q_j^\mu \left(\frac{\sin q_j (z^2 - x_j^2)^{1/2}}{(z^2 - x_j^2)^{1/2}} \right)^s,$$

$$\|f_j\|_{PW_+^{2,\nu}}^2 = 2 \int_0^1 (1-u^2)^\nu \left(\frac{|\sin u|}{u} \right)^{2s} u du + 2 \int_0^{+\infty} (1+u^2)^\nu \left(\frac{|\sin u|}{u} \right)^{2s} u du,$$

$$\|f_j\|_{PW_+^{2,\nu}}^2/c_6 \leq \sum_{k=1}^{\infty} (1 + |\operatorname{Im} \lambda_k|)(1 + |\lambda_k|)^{2\nu+1} e^{-2|\operatorname{Im} \lambda_k|} |f_j(\lambda_k)|^2 \leq c_6 \|f_j\|_{PW_+^{2,\nu}}^2,$$

and

$$\|f_j\|_{PW_+^{2,\nu}}^2/c_6 \leq \sum_{k=1}^{\infty} (1 + |\operatorname{Im} \lambda_k|)(1 + |\lambda_k|)^{2\nu+1} e^{-2|\operatorname{Im} \lambda_k|} |f_j(\lambda_k)|^2 \rightarrow 0, \quad j \rightarrow \infty.$$

We have a contradiction, since $\|f_j\|_{PW_+^{2,\nu}}^2$ is independent of j . \square

Lemma 3.7. *Let $\nu \geq -1/2$. If $P = \{\rho_k : k \in \mathbb{N}\}$ is a complete interpolating sequence for $PW_+^{2,\nu}$, then there exists a relatively dense set $\Gamma = \{\gamma_j : j \in \mathbb{Z} \setminus \{0\}\} \subset \Lambda$ such that $(|\gamma_j|^{2\nu+1} |S'(\gamma_j)|^2)$ satisfies the discrete (A_2) condition.*

Proof. Indeed, let $r > r_0$, $Q_j = Q(4jr; r)$ and $\Gamma = \{\gamma_j : j \in \mathbb{Z} \setminus \{0\}\} \subset \Lambda$ is a relatively dense sequence such that $\gamma_j \in Q_j$. Let $\Sigma = \{\sigma_j\}$ be another sequence with $|\gamma_j - \sigma_j| = \varepsilon$ and $S(\sigma_j) = \varepsilon S'(\gamma_j)$. Then the sequence $\{\sigma_j\}$ also satisfies condition (3.3). Therefore, by Lemma 3.5, we have

$$\sum_j |\sigma_j|^{2\nu+1} (1 + |\operatorname{Im} \sigma_j|) e^{-2|\operatorname{Im} \sigma_j|} |f(\sigma_j)|^2 \leq c_7 \|f\|_{PW_+^{2,\nu}}^2.$$

Since S' is an odd function, then for a finite set $\{d_j : j \in [1; m] \cap \mathbb{Z}\}$ the unique solution of the interpolation problem $f(\gamma_k) = d_k$, $\gamma_k \in \Gamma$ and $f(\gamma_k) = 0$, $\gamma_k \notin \Gamma$, has the form

$$f(z) = \sum_{k=1}^m \frac{2\gamma_k d_k S(z)}{(z^2 - \gamma_k^2) S'(\gamma_k)} = \sum_{k=-m, k \neq 0}^m \frac{d_k S(z)}{(z - \gamma_k) S'(\gamma_k)}, \quad d_{-k} := d_k.$$

In this case, according to Lemma 3.6, we get

$$c_6 \|f\|_{PW_+^{2,\nu}}^2 \leq \sum_j |\gamma_j|^{2\nu+1} (1 + |\operatorname{Im} \gamma_j|) e^{-2|\operatorname{Im} \gamma_j|} |d_j|^2.$$

Since $\gamma_j \in Q_j$, then the sequences $(|\operatorname{Im} \gamma_j|)$ and $(|\operatorname{Im} \sigma_j|)$ are bounded. Therefore

$$\sum_j |\sigma_j|^{2\nu+1} (1 + |\operatorname{Im} \sigma_j|) e^{-2|\operatorname{Im} \sigma_j|} |f(\sigma_j)|^2 \leq c_8 \sum_j |\gamma_j|^{2\nu+1} (1 + |\operatorname{Im} \gamma_j|) e^{-2|\operatorname{Im} \gamma_j|} |d_j|^2$$

and

$$\sum_j |\sigma_j|^{2\nu+1} |f(\sigma_j)|^2 \leq c_9 \sum_j |\gamma_j|^{2\nu+1} |d_j|^2.$$

Since

$$f(\sigma_j) = \varepsilon S'(\gamma_j) \sum_k \frac{d_k}{(\sigma_j - \gamma_k) S'(\gamma_k)},$$

we obtain

$$\sum_j |\gamma_j|^{2\nu+1} |S'(\gamma_j)|^2 \left| \sum_k \frac{d_k}{\sigma_j - \gamma_k} \right|^2 \leq c_9 \sum_j |\gamma_j|^{2\nu+1} |S'(\gamma_j)|^2 |d_j|^2.$$

Thus, the operator $\mathcal{H}_{\Gamma,\Sigma} : d = \{d_j\} \mapsto \mathcal{H}_{\Gamma,\Sigma} d$, $\mathcal{H}_{\Gamma,\Sigma} d := \sum_k \frac{d_k}{\sigma_j - \gamma_k}$ is bounded on $l_+^{2,w}$ if $w_j = |\gamma_j|^{2\nu+1} |S'(\gamma_j)|^2$. Therefore ([17]), the sequence $(|\gamma_j|^{2\nu+1} |S'(\gamma_j)|^2)$ satisfies the discrete (A_2) condition, and the lemma is proved. \square

Lemma 3.8. *If $\nu \geq -1/2$ and $P = \{\rho_k : k \in \mathbb{N}\}$ is a complete interpolating sequence for $PW_+^{2,\nu}$, then 5) holds.*

In fact, since the discrete (A_2) condition is equivalent to the continuous (A_2) condition, by using Lemma 2 from [17, p. 368] and the inequality $t^s s^{1-\alpha} \leq t + s$, $t, s > 0$, $\alpha \in [0; 1]$, we obtain that the statement of this lemma follows from Lemma 3.7.

Similarly to [17], by using conditions 1), 2) and 5), we obtain the following lemma.

Lemma 3.9. (see [17, Lemma 3, p. 371]) *Let condition (3.4) be true. Then*

$$|S(z)| \geq c_{10} (1 + |z|)^{-1/2} e^{|\operatorname{Im} z|}, \quad \text{for } \operatorname{dist}(z; \Lambda) \geq \varepsilon (1 + |\operatorname{Im} z|).$$

Lemma 3.10. *Let $\nu \geq -1/2$ and $P = \{\rho_k : k \in \mathbb{N}\}$ be an arbitrary sequence of nonzero complex numbers. A sequence $\Lambda = \{\lambda_k : k \in \mathbb{Z} \setminus \{0\}\}$ is a complete interpolating sequence for $PW_+^{2,\nu}$ if and only if conditions 1) – 5) hold.*

Proof. The necessity follows from Lemmas 3.3–3.8. Let us prove the sufficiency. First, we will show that the function

$$f(z) = \sum_{m \in \mathbb{N}} \frac{2\rho_m d_m S(z)}{(z^2 - \rho_m^2) S'(\lambda_m)} = v.p. \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{d_m S(z)}{(z - \lambda_m) S'(\lambda_m)} \quad (3.7)$$

is the required solution of the interpolation problem $f(\rho_k) = d_k$, where $d_{-k} := d_k$ for $k \in \mathbb{N}$. To this end, it suffices to estimate the partial sums of series (3.7) corresponding to $\lambda_m \in \mathbb{R} \cup \mathbb{C}_{0,+\infty}$ and $\lambda_m \in \mathbb{R} \cup \mathbb{C}_{-\infty,0}$, respectively on the lines $\operatorname{Im} z = -1/2$ and $\operatorname{Im} z = 1/2$. Following [17, p. 373], we will

give the corresponding estimates on the line $\text{Im } z = 0$ assuming that $\text{Im } \lambda_m \geq 1/2$ and $\text{Im } \lambda_m < 1/2$, respectively. In the first case, let

$$B(z) = \prod_{\text{Im } \lambda_j \geq 1/2} \frac{z - \lambda_j}{z - \bar{\lambda}_j}, \quad G(z) = \frac{S(z)}{e^{-iz} B(z)}.$$

Then $S(z) = G(z)e^{-iz}B(z)$, where G is a bounded outer function in \mathbb{C}^+ , and we observe that 5) is equivalent to $|G(x)|^2$ satisfying the (A_2) condition. Moreover, $|G(x)|^{-2}$ satisfies the (A_2) condition, and the Lemma 3.4 implies $|S'(\lambda_k)| \asymp |G(\lambda_k)| \frac{e^{\text{Im } \lambda_k}}{\text{Im } \lambda_k}$. Now let

$$\mathcal{H} : f \mapsto \mathcal{H}f(t) = \frac{1}{i\pi} \int_{-\infty}^{+\infty} \frac{f(\tau)}{t - \tau} d\tau$$

is the classical Hilbert operator. Following [17, p. 373], we consider the function

$$\tilde{f}(x) = \sum_{\substack{\text{Im } \lambda_m \geq 1/2, \\ m=-n, m \neq 0}}^n \frac{\text{Im } \lambda_m d_m e^{-\text{Im } \lambda_m} G(x)}{(x - \lambda_m) G(\lambda_m)}.$$

By duality (see [17, p. 373]), we obtain (here $w(x) = x^{\nu+1/2}$, $h_w(x) = w(x)h(x) \in L^2(\mathbb{R})$)

$$\begin{aligned} \|\tilde{f}\|_{L^{2,\nu}(\mathbb{R})} &= \sup_{\substack{\|h\| \leq 1, \\ h \in L^{2,\nu}(\mathbb{R})}} \left| \sum_{\substack{\text{Im } \lambda_m \geq 1/2, \\ m=-n, m \neq 0}}^n \int_{-\infty}^{+\infty} \frac{\text{Im } \lambda_m d_m e^{-\text{Im } \lambda_m} G(x) h_w(x)}{(x - \lambda_m) G(\lambda_m)} dx \right| \\ &\leq \sup_{\substack{\|h\| \leq 1, \\ h \in L^{2,\nu}(\mathbb{R})}} \left| \sum_{\substack{\text{Im } \lambda_m \geq 1/2, \\ m=-n, m \neq 0}}^n \frac{\text{Im } \lambda_m d_m e^{-\text{Im } \lambda_m} \mathcal{H}G h_w(\lambda_m)}{G(\lambda_m)} \right| \\ &\leq \sup_{\substack{\|h\| \leq 1, \\ h \in L^{2,\nu}(\mathbb{R})}} \left(\sum_{\text{Im } \lambda_m \geq 1/2} \left| \frac{\mathcal{H}G h_w(\lambda_m)}{G} \right|^2 \text{Im } \lambda_m \right)^{1/2} \\ &\quad \times \left(\sum_{\text{Im } \lambda_m \geq 1/2} \text{Im } \lambda_m e^{-2\text{Im } \lambda_m} |d_m|^2 \right)^{1/2}. \end{aligned}$$

Since $\sum_{\text{Im } \lambda_k \geq 0} \text{Im } \lambda_k \delta_{\lambda_k}$ is a Carleson measure, $|G(x)|^{-2}$ satisfies the (A_2) condition, G is an outer function in \mathbb{C}^+ , we have $\mathcal{H}Gh/G \in H^2(\mathbb{C}^+)$. Therefore, the last sums are uniformly bounded, and we get the desired conclusion. The sum corresponding to $\text{Im } \lambda_k < 0$ is treated similarly. Hence, there exists a solution of the considered interpolation problem. Now we turn to the proof of uniqueness. Observe first that (see [17, p. 370])

$$\int_{-\infty}^{+\infty} |F(x)|^2 \frac{dx}{1 + |x|^2} < +\infty, \quad \int_{-\infty}^{+\infty} |F(x)|^2 dx = +\infty.$$

Suppose that $f(\rho) = 0$, $\rho \in P$. Let $\psi(z) = f(z)/S(z)$. Since, by Lemma 3.2, $|f(z)| \leq c_4 \|f\|_{PW_+^{2,\nu}} e^{|\text{Im } z|} (1 + |z|)^{-\nu-1/2} (1 + |\text{Im } z|)^{-1/2}$, $z \in \mathbb{C}$, if $f \in PW_+^{2,\nu}$, then using Lemma 3.9, we obtain

$$|\psi(z)| = \left| \frac{f(z)}{S(z)} \right| \leq c_{11} \frac{e^{|\text{Im } z|} (1 + |z|)^{-\nu-1/2} (1 + |\text{Im } z|)^{-1/2}}{(1 + |z|)^{-1/2} e^{|\text{Im } z|}} = c_{11} \frac{(1 + |z|)^{-\nu}}{(1 + |\text{Im } z|)^{1/2}}.$$

Therefore, $|\psi(z)|$ is uniformly bounded for z satisfying $\text{dist}(z; \Lambda) \geq \varepsilon(1 + |\text{Im } z|)$. By the classical Phragmén-Lindelöf principle ([16, p. 39]), we get $\psi(z) \equiv c_{12}$, whereas ([17, p. 372]) $\int_{-\infty}^{+\infty} |S(x + i)|^2 dx = +\infty$ and $\int_{-\infty}^{+\infty} |S(x)|^2 dx = +\infty$. Hence, $\psi(z) \equiv 0$. \square

Theorem 1.1 is an immediate corollary of Lemma 3.10 and Theorem 1.2.

Acknowledgments

The authors thank the unknown Referee for a careful reading of this paper. The first version of the paper was significantly improved by the suggestions of the Referee.

References

- [1] N.I. Akhiezer, *To the theory of paired integral equations*, Uchenye Zapiski Kharkov Gos. Univ. 25 (1957), 5–31 (in Russian).
- [2] R.P. Boas, H. Pollard, *Complete sets of Bessel and Legendre functions*, Ann. of Math. 48 (1947), no. 2, 366–384.
- [3] M.M. Dzhrbashyan, *Integral transforms and representations of functions in the complex domain*, Nauka, Moscow, 1966 (in Russian).
- [4] M.M. Dzhrbashyan, S.G. Rafayelyan, *On entire functions of exponential type in weighted L^2 classes*, Dokl. Akad. Nauk Arm. SSR 73 (1981), no. 1, 29–36 (in Russian).
- [5] M.M. Dzhrbashyan, S.G. Rafayelyan, *Interpolation expansions in classes of entire functions and Riesz bases generated by them*, Izv. Akad. Nauk Arm. SSR. Matematika 22 (1987), no. 1, 23–63. English transl. in Sov. J. Contemp. Math. Anal., Arm. Acad. Sci. 22 (1987), no. 1, 20–61 (in Russian).
- [6] M.M. Djrbashian, S.G. Raphaelian, *Interpolation theorems and expansions with respect to Fourier type systems*, J. Approx. Theory 50 (1987), no. 4, 297–325.
- [7] J.B. Garnett, *Bounded analytic functions*, Academic Press, 1981.
- [8] J.L. Griffith, *Hankel transforms of functions zero outside a finite interval*, J. Proc. Roy. Soc. New South Wales 89 (1955), 109–115.
- [9] G.M. Gubreev, *The basis property of a family of Mittag-Leffler type functions, Djrbashian's transformation and weighted estimates of Cauchy type integrals*, Izv. Akad. Nauk Arm. SSR. Matematika 23 (1988), no. 3, 237–269 (in Russian). English transl. in Sov. J. Contemp. Math. Anal., Arm. Acad. Sci. 23 (1988), no. 3, 43–78.
- [10] G.M. Gubreev, *An integral transformation of Djrbashian type and interpolation by entire functions of finite order*, Izv. Akad. Nauk Arm. SSR. Matematika 25 (1990), no. 1, 83–90 (in Russian). English transl. in Sov. J. Contemp. Math. Anal., Arm. Acad. Sci. 25 (1990), no. 1, 79–85.
- [11] G.M. Gubreev, *Spectral analysis of biorthogonal expansions, generated by Muckenhoupt weights*, Zap. Nauchn. Sem. POMI 190 (1991), 34–80 (in Russian). English transl. in J. Math. Sci. (N.Y.) 71 (1994), no. 1, 2192–2221.
- [12] G.M. Gubreev, V.N. Levchuk, *Description of unconditional bases formed by values of the Dunkl kernels*, Funktsional. Anal. i Prilozhen 49 (2015), no. 1, 79–82 (in Russian). English transl. in Funct. Anal. Appl. 49 (2015), no. 1, 64–66.
- [13] H. Hochstadt, *The mean convergence of Fourier-Bessel series*, SIAM Rev. 9 (1967), 211–218.
- [14] S.V. Khrushchev, N.K. Nikol'skii, B.S. Pavlov, *Unconditional bases of exponentials and reproducing kernels*, in Complex Analysis and Spectral Theory (ed. V.P. Havin and N.K. Nikol'skii), Lecture Notes in Math. 864, Springer-Verlag (1981), 214–335.
- [15] P. Koosis, *Introduction to H_p spaces*, Second edition. Cambridge Tracts in Mathematics 115, Cambridge University Press, Cambridge, 1998.
- [16] B.Ya. Levin, *Lectures on entire functions*, Transl. Math. Monogr. 150. Amer. Math. Soc., Providence, R.I., 1996.
- [17] Yu.I. Lyubarskii, K. Seip, *Complete interpolating sequences for Paley-Wiener spaces and Muckenhoupt's (A_p) condition*, Rev. Mat. Iberoam 13 (1997), no. 2, 361–376.
- [18] Yu.I. Lyubarskii, K. Seip, *Weighted Paley-Wiener spaces*, J. Amer. Math. Soc. 15 (2002), no. 4, 979–1006.
- [19] A.M. Minkin, *Reflection of exponents and unconditional bases of exponentials*, Algebra i Analiz 3 (1991), no. 5, 109–134 (in Russian). English transl. in St. Petersburg Math. J. 3 (1992), no. 5, 1043–1064.
- [20] N.K. Nikol'skii, *Bases of exponentials and the values of reproducing kernels*, Dokl. Akad. Nauk SSSR 252 (1980), 1316–1320 (in Russian). English transl. in Sov. Math. Dokl. 21 (1980), 937–941.

- [21] B.S. Pavlov, *Basicity of an exponential system and Muckenhoupt's condition*, Dokl. Akad. Nauk SSSR 247 (1979), no. 1, 37–40 (in Russian). English transl. in Sov. Math. Dokl. 20 (1979), no. 4, 655–659.
- [22] S.G. Rafayelyan, *Interpolation and basisness in weighted classes of entire functions of exponential type*, Izv. Akad. Nauk Arm. SSR, Matematika 18 (1983), no. 3, 167–186 (in Russian). English transl. in Sov. J. Contemp. Math. Anal., Arm. Acad. Sci. 18 (1983), no. 3, 1–21.
- [23] S.G. Rafayelyan, *Basisness of some biorthogonal systems in $L^2(-\sigma; \sigma)$ with weights*, Izv. Akad. Nauk Arm. SSR, Matematika 19 (1984), no. 3, 207–218 (in Russian). English transl. in Sov. J. Contemp. Math. Anal., Arm. Acad. Sci. 19 (1984), no. 3, 21–32.
- [24] A.M. Sedletsii, *Analytic Fourier transforms and exponential approximations, II*, Sovrem. Mat. Fundam. Napravl. 6 (2003), 3–162 (in Russian). English transl. in J. Math. Sci. (N.Y.) 130 (2005), no. 6, 5083–5255.
- [25] I. Singer, *Bases in Banach spaces*, Vol. 1, Springer-Verlag, Berlin, 1970.
- [26] K. Stempak, *On convergence and divergence of Fourier-Bessel series*, Electron. Trans. Numer. Anal. 14 (2002), 223–235.
- [27] V.S. Vladimirov, *Equations of mathematical physics*, Nauka, Moscow, 1981 (in Russian). English transl. in Mir Publishers, Moscow, 1984.
- [28] B.V. Vynnyts'kyi, R.V. Khats', *Completeness and minimality of systems of Bessel functions*, Ufa Math. J. 5 (2013), no. 2, 131–141.
- [29] B.V. Vynnyts'kyi, R.V. Khats', *On the completeness and minimality of sets of Bessel functions in weighted L^2 -spaces*, Eurasian Math. J. 6 (2015), no. 1, 123–131.
- [30] B.V. Vynnyts'kyi, R.V. Khats', *A remark on basis property of systems of Bessel and Mittag-Leffler type functions*, Izv. NAN Armenii, Matematika 50 (2015), no. 6, 16–25 (in Russian). English transl. in J. Contemp. Math. Anal. 50 (2015), no. 6, 300–305.
- [31] B.V. Vynnyts'kyi, R.V. Khats', *Complete biorthogonal systems of Bessel functions*, Mat. Stud. 48 (2017), no. 2, 150–155.
- [32] B.V. Vynnyts'kyi, R.V. Khats', *Some approximation properties of the systems of Bessel functions of index $-3/2$* , Mat. Stud. 34 (2010), no. 2, 152–159.
- [33] G.N. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press, Cambridge, 1944.

Bohdan Vasil'evich Vynnyts'kyi, Ruslan Vasil'evich Khats', Iryna Bogdanovna Sheparovych
 Institute of Physics, Mathematics, Economy and Innovation Technologies
 Drohobych Ivan Franko State Pedagogical University
 3 Stryis'ka St., 82100 Drohobych, Ukraine
 E-mails: khats@ukr.net, isheparovych@ukr.net

Received: 27.03.2019