

ISSN (Print): 2077-9879  
ISSN (Online): 2617-2658

# Eurasian Mathematical Journal

2022, Volume 13, Number 2

Founded in 2010 by  
the L.N. Gumilyov Eurasian National University  
in cooperation with  
the M.V. Lomonosov Moscow State University  
the Peoples' Friendship University of Russia (RUDN University)  
the University of Padua

Starting with 2018 co-funded  
by the L.N. Gumilyov Eurasian National University  
and  
the Peoples' Friendship University of Russia (RUDN University)

Supported by the ISAAC  
(International Society for Analysis, its Applications and Computation)  
and  
by the Kazakhstan Mathematical Society

Published by  
the L.N. Gumilyov Eurasian National University  
Nur-Sultan, Kazakhstan

# EURASIAN MATHEMATICAL JOURNAL

## Editorial Board

### Editors-in-Chief

V.I. Burenkov, M. Otelbaev, V.A. Sadovnichy

### Vice-Editors-in-Chief

K.N. Ospanov, T.V. Tararykova

### Editors

Sh.A. Alimov (Uzbekistan), H. Begehr (Germany), T. Bekjan (Kazakhstan), O.V. Besov (Russia), N.K. Blied (Kazakhstan), N.A. Bokayev (Kazakhstan), A.A. Borubaev (Kyrgyzstan), G. Bourdaud (France), A. Caetano (Portugal), M. Carro (Spain), A.D.R. Choudary (Pakistan), V.N. Chubarikov (Russia), A.S. Dzumadildaev (Kazakhstan), V.M. Filippov (Russia), H. Ghazaryan (Armenia), M.L. Goldman (Russia), V. Goldshtein (Israel), V. Guliyev (Azerbaijan), D.D. Haroske (Germany), A. Hasanoglu (Turkey), M. Huxley (Great Britain), P. Jain (India), T.Sh. Kalmenov (Kazakhstan), B.E. Kangyzhin (Kazakhstan), K.K. Kenzhibayev (Kazakhstan), S.N. Kharin (Kazakhstan), E. Kissin (Great Britain), V. Kokilashvili (Georgia), V.I. Korzyuk (Belarus), A. Kufner (Czech Republic), L.K. Kussainova (Kazakhstan), P.D. Lamberti (Italy), M. Lanza de Cristoforis (Italy), F. Lanzara (Italy), V.G. Maz'ya (Sweden), K.T. Mynbayev (Kazakhstan), E.D. Nursultanov (Kazakhstan), R. Oinarov (Kazakhstan), I.N. Parasidis (Greece), J. Pečarić (Croatia), S.A. Plaksa (Ukraine), L.-E. Persson (Sweden), E.L. Presman (Russia), M.A. Ragusa (Italy), M.D. Ramazanov (Russia), M. Reissig (Germany), M. Ruzhansky (Great Britain), M.A. Sadybekov (Kazakhstan), S. Sagitov (Sweden), T.O. Shaposhnikova (Sweden), A.A. Shkalikov (Russia), V.A. Skvortsov (Poland), G. Sinamon (Canada), E.S. Smailov (Kazakhstan), V.D. Stepanov (Russia), Ya.T. Sultanaev (Russia), D. Suragan (Kazakhstan), I.A. Taimanov (Russia), J.A. Tussupov (Kazakhstan), U.U. Umirbaev (Kazakhstan), Z.D. Usmanov (Tajikistan), N. Vasilevski (Mexico), Dachun Yang (China), B.T. Zhumagulov (Kazakhstan)

### Managing Editor

A.M. Temirkhanova

## Aims and Scope

The Eurasian Mathematical Journal (EMJ) publishes carefully selected original research papers in all areas of mathematics written by mathematicians, principally from Europe and Asia. However papers by mathematicians from other continents are also welcome.

From time to time the EMJ publishes survey papers.

The EMJ publishes 4 issues in a year.

The language of the paper must be English only.

The contents of the EMJ are indexed in Scopus, Web of Science (ESCI), Mathematical Reviews, MathSciNet, Zentralblatt Math (ZMATH), Referativnyi Zhurnal – Matematika, Math-Net.Ru.

The EMJ is included in the list of journals recommended by the Committee for Control of Education and Science (Ministry of Education and Science of the Republic of Kazakhstan) and in the list of journals recommended by the Higher Attestation Commission (Ministry of Education and Science of the Russian Federation).

## Information for the Authors

Submission. Manuscripts should be written in LaTeX and should be submitted electronically in DVI, PostScript or PDF format to the EMJ Editorial Office through the provided web interface ([www.enu.kz](http://www.enu.kz)).

When the paper is accepted, the authors will be asked to send the tex-file of the paper to the Editorial Office.

The author who submitted an article for publication will be considered as a corresponding author. Authors may nominate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not guaranteed.

Copyright. When the paper is accepted, the copyright is automatically transferred to the EMJ. Manuscripts are accepted for review on the understanding that the same work has not been already published (except in the form of an abstract), that it is not under consideration for publication elsewhere, and that it has been approved by all authors.

Title page. The title page should start with the title of the paper and authors' names (no degrees). It should contain the Keywords (no more than 10), the Subject Classification (AMS Mathematics Subject Classification (2010) with primary (and secondary) subject classification codes), and the Abstract (no more than 150 words with minimal use of mathematical symbols).

Figures. Figures should be prepared in a digital form which is suitable for direct reproduction.

References. Bibliographical references should be listed alphabetically at the end of the article. The authors should consult the Mathematical Reviews for the standard abbreviations of journals' names.

Authors' data. The authors' affiliations, addresses and e-mail addresses should be placed after the References.

Proofs. The authors will receive proofs only once. The late return of proofs may result in the paper being published in a later issue.

Offprints. The authors will receive offprints in electronic form.

## Publication Ethics and Publication Malpractice

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the EMJ implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The EMJ follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (<http://publicationethics.org/files/u2/NewCode.pdf>). To verify originality, your article may be checked by the originality detection service CrossCheck <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the EMJ.

The Editorial Board of the EMJ will monitor and safeguard publishing ethics.

# The procedure of reviewing a manuscript, established by the Editorial Board of the Eurasian Mathematical Journal

## 1. Reviewing procedure

1.1. All research papers received by the Eurasian Mathematical Journal (EMJ) are subject to mandatory reviewing.

1.2. The Managing Editor of the journal determines whether a paper fits to the scope of the EMJ and satisfies the rules of writing papers for the EMJ, and directs it for a preliminary review to one of the Editors-in-chief who checks the scientific content of the manuscript and assigns a specialist for reviewing the manuscript.

1.3. Reviewers of manuscripts are selected from highly qualified scientists and specialists of the L.N. Gumilyov Eurasian National University (doctors of sciences, professors), other universities of the Republic of Kazakhstan and foreign countries. An author of a paper cannot be its reviewer.

1.4. Duration of reviewing in each case is determined by the Managing Editor aiming at creating conditions for the most rapid publication of the paper.

1.5. Reviewing is confidential. Information about a reviewer is anonymous to the authors and is available only for the Editorial Board and the Control Committee in the Field of Education and Science of the Ministry of Education and Science of the Republic of Kazakhstan (CCFES). The author has the right to read the text of the review.

1.6. If required, the review is sent to the author by e-mail.

1.7. A positive review is not a sufficient basis for publication of the paper.

1.8. If a reviewer overall approves the paper, but has observations, the review is confidentially sent to the author. A revised version of the paper in which the comments of the reviewer are taken into account is sent to the same reviewer for additional reviewing.

1.9. In the case of a negative review the text of the review is confidentially sent to the author.

1.10. If the author sends a well reasoned response to the comments of the reviewer, the paper should be considered by a commission, consisting of three members of the Editorial Board.

1.11. The final decision on publication of the paper is made by the Editorial Board and is recorded in the minutes of the meeting of the Editorial Board.

1.12. After the paper is accepted for publication by the Editorial Board the Managing Editor informs the author about this and about the date of publication.

1.13. Originals reviews are stored in the Editorial Office for three years from the date of publication and are provided on request of the CCFES.

1.14. No fee for reviewing papers will be charged.

## 2. Requirements for the content of a review

2.1. In the title of a review there should be indicated the author(s) and the title of a paper.

2.2. A review should include a qualified analysis of the material of a paper, objective assessment and reasoned recommendations.

2.3. A review should cover the following topics:

- compliance of the paper with the scope of the EMJ;
- compliance of the title of the paper to its content;
- compliance of the paper to the rules of writing papers for the EMJ (abstract, key words and phrases, bibliography etc.);
- a general description and assessment of the content of the paper (subject, focus, actuality of the topic, importance and actuality of the obtained results, possible applications);
- content of the paper (the originality of the material, survey of previously published studies on the topic of the paper, erroneous statements (if any), controversial issues (if any), and so on);

- exposition of the paper (clarity, conciseness, completeness of proofs, completeness of bibliographic references, typographical quality of the text);
- possibility of reducing the volume of the paper, without harming the content and understanding of the presented scientific results;
- description of positive aspects of the paper, as well as of drawbacks, recommendations for corrections and complements to the text.

2.4. The final part of the review should contain an overall opinion of a reviewer on the paper and a clear recommendation on whether the paper can be published in the Eurasian Mathematical Journal, should be sent back to the author for revision or cannot be published.

## Web-page

The web-page of the EMJ is [www.emj.enu.kz](http://www.emj.enu.kz). One can enter the web-page by typing Eurasian Mathematical Journal in any search engine (Google, Yandex, etc.). The archive of the web-page contains all papers published in the EMJ (free access).

## Subscription

Subscription index of the EMJ 76090 via KAZPOST.

## E-mail

[eurasianmj@yandex.kz](mailto:eurasianmj@yandex.kz)

The Eurasian Mathematical Journal (EMJ)  
The Nur-Sultan Editorial Office  
The L.N. Gumilyov Eurasian National University  
Building no. 3  
Room 306a  
Tel.: +7-7172-709500 extension 33312  
13 Kazhymukan St  
010008 Nur-Sultan, Kazakhstan

The Moscow Editorial Office  
The Peoples' Friendship University of Russia  
(RUDN University)  
Room 473  
3 Ordzonikidze St  
117198 Moscow, Russia

**A MULTI-POINT PROBLEM FOR A SYSTEM OF DIFFERENTIAL EQUATIONS WITH PIECEWISE-CONSTANT ARGUMENT OF GENERALIZED TYPE AS A NEURAL NETWORK MODEL****A. Abildayeva, A. Assanova, A. Imanchiyev**

Communicated by K.N. Ospanov

**Key words:** differential equations with piecewise-constant argument of generalized type, neural network model, multi-point boundary value problem, solvability criteria, algorithms of parameterization method.

**AMS Mathematics Subject Classification:** 34A36, 34K10.

**Abstract.** We consider a system of ordinary differential equations with piecewise-constant argument of generalized type. An interval is divided into  $N$  parts, the values of a solution at the interior points of the subintervals are considered as additional parameters, and a system of ordinary differential equations with piecewise-constant argument of generalized type is reduced to the Cauchy problems on the subintervals for linear system of ordinary differential equations with parameters. Using the solutions to these problems, new general solutions to system of differential equations with piecewise-constant argument of generalized type are introduced and their properties are established. Based on the general solution, boundary condition, and continuity conditions of a solution at the interior points of the partition, the system of linear algebraic equations with respect to parameters is composed. Its coefficients and right-hand sides are found by solving the Cauchy problems for a linear system of ordinary differential equations on the subintervals. It is shown that the solvability of boundary value problems is equivalent to the solvability of composed systems. Methods for solving boundary value problems are proposed, which are based on the construction and solving of these systems.

**DOI:** <https://doi.org/10.32523/2077-9879-2022-13-2-08-17>

## 1 Introduction and statement of problem

It is well known that mathematical modeling of processes with discontinuity effects has necessitated the need to develop the theory of differential equations with discontinuities. An important class of such equations is comprised of differential equations with a piecewise constant argument (DEPCA). The study of DEPCA was initiated by Busenberg, Cooke, Shah, and Wiener [22], [19], [39]. The problems of the existence and uniqueness of solutions to DEPCA, their oscillations and stability, integral manifolds and periodic solutions have been extensively discussed by many authors [33], [20], [34], [40], [23], [38], [15], [16], [14].

When modeling DEPCA, the deviation of the argument, taken as the greatest integer function, is always constant and equal to one. But this approach can contradict real phenomena. The generalization of DEPCA has been undertaken by M.U.Akhmet [1], [2], [3], [4]. In his works the greatest integer function as deviating argument was replaced by an arbitrary piecewise constant function. Thus, differential equations with piecewise constant argument of generalized type (DEPCAG) are more suitable for modeling and solving various applied problems, including areas of neural networks,



discontinuous dynamical systems, hybrid systems, etc. To date, the theory of DEPCAG on the entire axis has been developed and their applications have been implemented. The results have been extended to periodic impulse systems of DEPCAG [5], [6], [11], [9], [10], [21], [7]. Note that an electronic neural networks were modeled as differential equations with piecewise constant arguments of generalized type [18], [11], [9]. By reducing these equations to an equivalent integral equation, some new stability conditions are obtained.

Along with the study of various properties of DEPCA, a number of authors investigated the problems of solvability and construction of solutions to boundary value problems for these equations on a finite interval [35], [37], [24], [8].

For DEPCAG, however, the problems of solvability of boundary value problems on a finite interval still remain open.

This issue can be resolved by developing constructive methods.

So, on  $[0, T]$ , we consider the following multi-point boundary value problem for a system of DEPCAG:

$$\frac{dx}{dt} = A(t)x + A_0(t)x(\gamma(t)) + f(t), \quad x \in \mathbb{R}^n, \quad t \in (0, T), \quad (1.1)$$

$$\sum_{i=0}^N B_i x(\theta_i) = d, \quad d \in \mathbb{R}^n. \quad (1.2)$$

Here  $x(t) = \text{col}(x_1(t), x_2(t), \dots, x_n(t))$  is the unknown function,  $(n \times n)$  matrices  $A(t)$ ,  $A_0(t)$  and  $n$ -vector  $f(t)$  are continuous on  $[0, T]$ ;

$\gamma(t) = \zeta_j$  if  $t \in [\theta_j, \theta_{j+1})$ ,  $j = \overline{0, N-1}$ ;  $\theta_j \leq \zeta_j \leq \theta_{j+1}$  for all  $j = \overline{0, 1, \dots, N-1}$ ;  $0 = \theta_0 < \theta_1 < \dots < \theta_{N-1} < \theta_N = T$ ;  $B_i$  are constant  $(n \times n)$  matrices,  $i = \overline{0, N}$ , and  $d$  is a constant vector.

The aim of the present paper is to develop a constructive method for investigating and solving the boundary value problem, including an algorithm for finding a solution to problem (1.1), (1.2) as well.

To this end, we use a new concept of general solution and Dzhumabaev's parametrization method [25], [26]. This concept of general solution has been introduced for the linear Fredholm integro-differential equation in [27] and for the linear loaded differential equation and a family of such equations in [28], [29]. New general solutions are also introduced to ordinary differential equations and their properties are established in [30]. Results are developed to nonlinear Fredholm integro-differential equations [31], [32] and to problems with a parameter for integro-differential equations [13]. Based on the general solution methods for solving boundary value problems are proposed.

The paper is organized as follows.

The interval  $[0, T]$  is divided into  $N$  parts according to the partition  $\Delta_N : \theta_0 = 0 < \theta_1 < \theta_2 < \dots < \theta_N = T$ , and the  $\Delta_N$  general solution to a linear system of differential equation with a piecewise-constant argument of generalized type is introduced. The  $\Delta_N$  general solution, denoted by  $x(\Delta_N, t, \lambda)$ , contains an arbitrary vectors  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_N) \in \mathbb{R}^{nN}$ . Using  $x(\Delta_N, t, \lambda)$ , we establish solvability criteria of considered problem and propose an algorithm for finding its solution.

A function  $x^*(t) : [0, T] \rightarrow \mathbb{R}^n$  is a solution to problem (1.1), (1.2) if:

- (i)  $x^*(t)$  is continuous on  $[0, T]$ ;
- (ii)  $x^*(t)$  is differentiable on  $[0, T]$  with the possible exception of the points  $\theta_j$ ,  $j = \overline{0, N-1}$ , at which the one-sided derivatives exist;
- (iii)  $x^*(t)$  satisfies the system of equations (1.1) on each interval  $(\theta_j, \theta_{j+1})$ ,  $j = \overline{0, N-1}$ ; at the points  $\theta_j$ ,  $j = \overline{0, N-1}$ , system (1) is satisfied by the right-hand derivative of  $x^*(t)$ ;
- (iv)  $x^*(t)$  satisfies boundary condition (1.2) at  $t = \theta_i$ ,  $i = \overline{0, N}$ .

## 2 Scheme of the method and $\Delta_N$ general solution

Let  $\Delta_N$  denote the partition of the interval  $[0, T)$  by points  $t = \theta_r$ ,  $r = \overline{1, N-1}$ :  
 $[0, T) = \bigcup_{r=1}^N [\theta_{r-1}, \theta_r)$ .

We define the following spaces:

$C([0, T], \mathbb{R}^n)$  is the space of all continuous functions  $x : [0, T] \rightarrow \mathbb{R}^n$  with the norm

$$\|x\|_1 = \max_{t \in [0, T]} \|x(t)\| = \max_{t \in [0, T]} \max_{i=1, n} |x_i(t)|;$$

$C([0, T], \Delta_N, \mathbb{R}^{nN})$  is the space of function systems  $x[t] = (x_1(t), x_2(t), \dots, x_N(t))$ , where  $x_r : [\theta_{r-1}, \theta_r) \rightarrow \mathbb{R}^n$  are continuous functions that have finite left-hand limits  $\lim_{t \rightarrow \theta_r - 0} x_r(t)$  for all  $r = \overline{1, N}$ , with the norm

$$\|x[\cdot]\|_2 = \max_{r=1, N} \sup_{t \in [\theta_{r-1}, \theta_r)} |x_r(t)|.$$

Denote by  $x_r(t)$  the restriction of a function  $x(t)$  to the  $r$ th interval  $[\theta_{r-1}, \theta_r)$ , i.e.

$$x_r(t) = x(t) \text{ for } t \in [\theta_{r-1}, \theta_r), \quad r = \overline{1, N}.$$

Then the function system  $x[t] = (x_1(t), x_2(t), \dots, x_N(t))$  belongs to  $C([0, T], \Delta_N, \mathbb{R}^{nN})$ , and its elements  $x_r(t)$ ,  $r = \overline{1, N}$ , satisfy the following system of ordinary differential equations with piecewise-constant argument of generalized type

$$\frac{dx_r}{dt} = A(t)x_r(t) + A_0(t)x_r(\zeta_{r-1}) + f(t), \quad t \in [\theta_{r-1}, \theta_r), \quad r = \overline{1, N}. \quad (2.1)$$

In (2.1) we take into account that  $\gamma(t) = \zeta_j$  if  $t \in [\theta_j, \theta_{j+1})$ ,  $j = \overline{0, N-1}$ .

We introduce additional parameters  $\lambda_r = x_r(\zeta_{r-1})$  for all  $r = \overline{1, N}$ . Making the substitution  $z_r(t) = x_r(t) - \lambda_r$  on every  $r$ -th interval  $[\theta_{r-1}, \theta_r)$ , we obtain the system of ordinary differential equations with parameters

$$\frac{dz_r}{dt} = A(t)(z_r(t) + \lambda_r) + A_0(t)\lambda_r + f(t), \quad t \in [\theta_{r-1}, \theta_r), \quad r = \overline{1, N}, \quad (2.2)$$

and initial conditions

$$z_r(\zeta_{r-1}) = 0, \quad r = \overline{1, N}. \quad (2.3)$$

Problems (2.2), (2.3) are Cauchy problems for system of ordinary differential equations with parameters on the intervals  $[\theta_{r-1}, \theta_r)$ ,  $r = \overline{1, N}$ . For any fixed  $\lambda_r \in \mathbb{R}^n$  and  $r$ , the Cauchy problem (2.2), (2.3) has a unique solution  $z_r(t, \lambda_r)$ , and the function system  $z[t, \lambda] = (z_1(t, \lambda_1), z_2(t, \lambda_2), \dots, z_N(t, \lambda_N))$  belongs to  $C([0, T], \Delta_N, \mathbb{R}^{nN})$ .

The function system  $z[t, \lambda]$  is referred to as a solution to Cauchy problems with parameters (2.2), (2.3). If a function system  $\tilde{x}[t] = (\tilde{x}_1(t), \tilde{x}_2(t), \dots, \tilde{x}_N(t))$  belongs to  $C([0, T], \Delta_N, \mathbb{R}^{nN})$ , and the functions  $\tilde{x}_r(t)$ ,  $r = \overline{1, N}$ , satisfy equations (2.1), then the function system  $z[t, \tilde{\lambda}] = (z_1(t, \tilde{\lambda}_1), z_2(t, \tilde{\lambda}_2), \dots, z_N(t, \tilde{\lambda}_N))$  with the elements  $z_r(t, \tilde{\lambda}_r) = \tilde{x}_r(t) - \tilde{\lambda}_r$ ,  $\tilde{\lambda}_r = \tilde{x}_r(\zeta_{r-1})$ ,  $r = \overline{1, N}$ , is a solution to the Cauchy problems with parameters (2.2), (2.3) for  $\lambda_r = \tilde{\lambda}_r$ ,  $r = \overline{1, N}$ . Conversely, if a function system  $z[t, \lambda^*] = (z_1(t, \lambda_1^*), z_2(t, \lambda_2^*), \dots, z_N(t, \lambda_N^*))$  is a solution to problems (2.2), (2.3) for  $\lambda_r = \lambda_r^*$ ,  $r = \overline{1, N}$ , then the function system  $x^*[t] = (x_1^*(t), x_2^*(t), \dots, x_N^*(t))$  with  $x_r^*(t) = \lambda_r^* + z_r(t, \lambda_r^*)$ ,  $r = \overline{1, N}$ , belongs to  $C([0, T], \Delta_N, \mathbb{R}^{nN})$ , and the functions  $x_r^*(t)$ ,  $r = \overline{1, N}$ , satisfy system of equations (2.1).

Let us now introduce a new general solution to the system of ordinary differential equations with piecewise-constant argument of generalized type (2.1).

**Definition 1.** Let  $z[t, \lambda] = (z_1(t, \lambda_1), z_2(t, \lambda_2), \dots, z_N(t, \lambda_N))$  be the solution to the Cauchy problems (2.2), (2.3) for the parameters  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_N) \in \mathbb{R}^{nN}$ . Then the function  $x(\Delta_N, t, \lambda)$ , given by the equalities

$$\begin{aligned} x(\Delta_N, t, \lambda) &= \lambda_r + z_r(t, \lambda_r), \text{ for } t \in [\theta_{r-1}, \theta_r], \quad r = \overline{1, N}, \text{ and} \\ x(\Delta_N, T, \lambda) &= \lambda_N + \lim_{t \rightarrow T-0} z_N(t, \lambda_N), \end{aligned}$$

is called the  $\Delta_N$  general solution to system of equations (2.1).

As follows from Definition 2.1, the  $\Delta_N$  general solution depends on  $N$  arbitrary vectors  $\lambda_r \in \mathbb{R}^n$  and satisfies system of equations (2.1) for all  $t \in (0, T) \setminus \{\theta_p, p = \overline{1, N-1}\}$ .

Take  $X_r(t)$ , a fundamental matrix of the ordinary differential equation

$$\frac{dz_r}{dt} = A(t)z_r(t), \quad t \in [\theta_{r-1}, \theta_r], \quad r = \overline{1, N},$$

and write down the solutions to the Cauchy problems with parameters (2.2), (2.3) in the form:

$$\begin{aligned} z_r(t) &= X_r(t) \int_{\zeta_{r-1}}^t X_r^{-1}(\tau)[A(\tau) + A_0(\tau)]d\tau \lambda_r + X_r(t) \int_{\zeta_{r-1}}^t X_r^{-1}(\tau)f(\tau)d\tau, \\ t &\in [\theta_{r-1}, \theta_r), \quad r = \overline{1, N}. \end{aligned}$$

Consider the Cauchy problems on the subintervals

$$\frac{dx}{dt} = A(t)x + P(t), \quad x(\zeta_{r-1}) = 0, \quad t \in [\theta_{r-1}, \theta_r], \quad r = \overline{1, N}, \quad (2.4)$$

where  $P(t)$  is a square matrix or a vector of dimension 2, continuous on  $[0, T]$ ,  $\theta_{r-1} \leq \zeta_{r-1} \leq \theta_r$  for all  $r = 1, 2, \dots, N$ . Denote by  $A_r(P, t)$  a unique solution to Cauchy problem (2.4) on each  $r$ th interval. The uniqueness of the solution to the Cauchy problem for linear ordinary differential equations yields

$$A_r(P, t) = X_r(t) \int_{\zeta_{r-1}}^t X_r^{-1}(\tau)P(\tau)d\tau, \quad t \in [\theta_{r-1}, \theta_r], \quad r = \overline{1, N}.$$

Therefore, we can represent the  $\Delta_N$  general solution to system of equations (2.1) in the form:

$$x(\Delta_N, t, \lambda) = \lambda_p + A_p(A + A_0, t)\lambda_p + A_p(f, t), \quad t \in [\theta_{p-1}, \theta_p], \quad p = \overline{1, N-1}, \quad (2.5)$$

$$x(\Delta_N, t, \lambda) = \lambda_N + A_N(A + A_0, t)\lambda_N + A_N(f, t), \quad t \in [\theta_{N-1}, \theta_N]. \quad (2.6)$$

The following statement affirms the function  $x(\Delta_N, t, \lambda)$  as a "general solution".

**Theorem 2.1.** *Let a piecewise continuous on  $[0, T]$  function  $\tilde{x}(t)$  with the possible discontinuity points  $t = \theta_p$ ,  $p = \overline{1, N-1}$ , be given, and  $x(\Delta_N, t, \lambda)$  be the  $\Delta_N$  general solution to system of equations (2.1). Suppose that the function  $\tilde{x}(t)$  has a continuous derivative and satisfies system of equations (2.1) for all  $t \in (0, T) \setminus \{\theta_p, p = \overline{1, N-1}\}$ . Then there exists a unique  $\tilde{\lambda} = (\tilde{\lambda}_1, \tilde{\lambda}_2, \dots, \tilde{\lambda}_N) \in \mathbb{R}^{nN}$  such that the equality  $x(\Delta_N, t, \tilde{\lambda}) = \tilde{x}(t)$  holds for all  $t \in [0, T]$ .*

The proof of this theorem is quite simple. Therefore, we do not present it.

**Corollary 2.1.** *Let  $x^*(t)$  be a solution to system of equations (2.1) and  $x(\Delta_N, t, \lambda)$  be the  $\Delta_N$  general solution to system of equations (2.1). Then there exists a unique  $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_N^*) \in \mathbb{R}^{nN}$  such that the equality  $x(\Delta_N, t, \lambda^*) = x^*(t)$  holds for all  $t \in [0, T]$ .*

If  $x(t)$  is a solution to system of equations (2.1), and  $x[t] = (x_1(t), x_2(t), \dots, x_N(t))$  is a function system composed of its restrictions to the subintervals  $[\theta_{r-1}, \theta_r]$ ,  $r = \overline{1, N}$ , then the equations

$$\lim_{t \rightarrow \theta_p - 0} x_p(t) = x_{p+1}(\theta_p), \quad p = \overline{1, N-1}, \quad (2.7)$$

hold. These equations are the continuity conditions for the solution to system of equations (2.1) at the interior points of the partition  $\Delta_N$ .

**Theorem 2.2.** *Let a function system  $x[t] = (x_1(t), x_2(t), \dots, x_N(t))$  belong to  $C([0, T], \Delta_N, \mathbb{R}^{nN})$ . Assume that the functions  $x_r(t)$ ,  $r = \overline{1, N}$ , satisfy system of equations (2.1) and continuity conditions (2.7). Then the function  $x^*(t)$ , given by the equalities*

$$\begin{aligned} x^*(t) &= x_r(t) \text{ for } t \in [\theta_{r-1}, \theta_r], \quad r = \overline{1, N}, \\ \text{and } x^*(T) &= \lim_{t \rightarrow T-0} x_N(t), \end{aligned}$$

*is continuous on  $[0, T]$ , continuously differentiable on  $(0, T)$  and satisfies system of equations (2.1).*

*Proof.* Equations (2.7), the equality  $x^*(T) = \lim_{t \rightarrow T-0} x_N(t)$ , and belonging of  $x[t] = (x_1(t), x_2(t), \dots, x_N(t))$  to  $C([0, T], \Delta_N, \mathbb{R}^{nN})$  ensure continuity of the function  $x^*(t)$  on the interval  $[0, T]$ . Since the functions  $x_r(t)$ ,  $r = \overline{1, N}$ , satisfy system of equations (3), the function  $x^*(t)$  has continuous derivative and satisfies system of equations (2.1) for all  $t \in [0, T] \setminus \{\theta_p, p = \overline{1, N-1}\}$ . The existence and continuity of the derivative of the function  $x^*(t)$  at the points  $t = \theta_p$ ,  $p = \overline{1, N-1}$ , follow from the relations:

$$\lim_{t \rightarrow \theta_p - 0} \dot{x}^*(t) = A(\theta_p)x^*(\theta_p) + A_0(\theta_p)x^*(\zeta_{p-1}) + f(\theta_p) = \lim_{t \rightarrow \theta_p + 0} \dot{x}^*(t), \quad p = \overline{1, N-1}.$$

Hence the function  $x^*(t)$  satisfies system of equations (2.1) at the interior points of the partition  $\Delta_N$  as well. □

### 3 Main results and algorithm

The  $\Delta_N$  general solution allows us to transfer the solvability of a multi-point boundary value problem to the solvability of a system of linear algebraic equations with respect to arbitrary vectors  $\lambda_r \in \mathbb{R}^2$ ,  $r = \overline{1, N}$ .

Substituting the suitable expressions of  $\Delta_N$  general solution (2.5), (2.6) into the multi-point condition (1.2) and continuity conditions (2.7), we obtain the system of linear algebraic equations

$$\sum_{i=0}^N B_i \left\{ I + A_i(A + A_0, \theta_i) \right\} \lambda_i = d - \sum_{i=0}^N B_i A_i(f, \theta_i), \quad (3.1)$$

$$\begin{aligned} &\left\{ I + A_p(A, \theta_p) \right\} \lambda_p - \left\{ I + A_{p+1}(A + A_0, \theta_p) \right\} \lambda_{p+1} \\ &= -A_p(f, \theta_p) + A_{p+1}(f, \theta_p), \quad p = \overline{1, N-1}, \end{aligned} \quad (3.2)$$

where  $I$  is the unit matrix of dimension  $n$ .

Denote by  $Q_*(\Delta_N)$   $nN \times nN$  matrix corresponding to the left-hand side of system (3.1), (3.2) and write the system as

$$Q_*(\Delta_N)\lambda = -F_*(\Delta_N), \quad \lambda \in \mathbb{R}^{nN}, \quad (3.3)$$

where  $F_*(\Delta_N) = \left( -d + \sum_{i=0}^N B_i A_i(f, \theta_i), A_1(f, \theta_1) - A_2(f, \theta_1), \right.$

$$\left. A_2(f, \theta_2) + A_3(f, \theta_2), \dots, A_{N-1}(f, \theta_{N-1}) + A_N(f, \theta_{N-1}) \right) \in \mathbb{R}^{nN}.$$

For any partition  $\Delta_N$ , Theorems 2.1 and 2.2 ensure the validity of the next assertion.

**Lemma 3.1.** *If  $x^*(t)$  is a solution to multi-point problem (1.1), (1.2) and  $\lambda_r^* = x^*(\zeta_{r-1})$ ,  $r = \overline{1, N}$ , then the vector  $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_N^*) \in \mathbb{R}^{nN}$  is a solution to system (3.3). Conversely, if  $\tilde{\lambda} = (\tilde{\lambda}_1, \tilde{\lambda}_2, \dots, \tilde{\lambda}_N) \in \mathbb{R}^{nN}$  is a solution to system (3.3) and  $z[t, \tilde{\lambda}] = (z_1(t, \tilde{\lambda}_1), z_2(t, \tilde{\lambda}_2), \dots, z_N(t, \tilde{\lambda}_N))$  is the solution to Cauchy problems (2.2), (2.3) for the parameter  $\tilde{\lambda} \in \mathbb{R}^{nN}$ , then the function  $\tilde{x}(t)$  given by the equalities  $\tilde{x}(t) = \tilde{\lambda}_r + z_r(t, \tilde{\lambda}_r)$ ,  $t \in [\theta_{r-1}, \theta_r]$ ,  $r = \overline{1, N}$ , and  $\tilde{x}(T) = \tilde{\lambda}_N + \lim_{t \rightarrow T-0} z_N(t, \tilde{\lambda}_N)$ , is a solution to multi-point problem (1.1), (1.2).*

**Definition 2.** The multi-point boundary value problem (1.1), (1.2) is called uniquely solvable if for any pair  $(f(t), d)$ , with  $f(t) \in C([0, T], \mathbb{R}^n)$  and  $d \in \mathbb{R}^n$ , it has a unique solution.

Lemma 3.1 and well-known theorems of linear algebra imply the following two statements.

**Theorem 3.1.** *The multi-point boundary value problem (1.1), (1.2) is solvable if and only if the vector  $F_*(\Delta_N)$  is orthogonal to the kernel of the transposed matrix  $(Q_*(\Delta_N))'$ , i.e. if and only if the equality*

$$(F_*(\Delta_N), \eta) = 0$$

*is valid for all  $\eta \in Ker(Q_*(\Delta_N))'$ , where  $(\cdot, \cdot)$  is the inner product in  $\mathbb{R}^{2N}$ .*

**Theorem 3.2.** *The multi-point boundary value problem (1.1), (1.2) is uniquely solvable if and only if  $nN \times nN$  matrix  $Q_*(\Delta_N)$  is invertible.*

So, by Theorems 3.1 and 3.2 it follows that the solvability of multi-point boundary value problem (1.1), (1.2) is equivalent to the solvability of system of algebraic equations (3.3). This system composed by solutions of Cauchy problems (2.2), (2.3), of multi-point condition (1.2) and continuity condition (2.7).

Based on the results of Section 3, we offer the following algorithm for finding a solution to the linear multi-point boundary value problem (1.1), (1.2).

**Algorithm.**

*Step 1.* Solve the Cauchy problems on the subintervals

$$\frac{dz}{dt} = A(t)z + A(t) + A_0(t), \quad z(\zeta_{r-1}) = 0, \quad t \in [\theta_{r-1}, \theta_r],$$

$$\frac{dz}{dt} = A(t)z + f(t), \quad z(\zeta_{r-1}) = 0, \quad t \in [\theta_{r-1}, \theta_r],$$

and find  $A_r(A + A_0, \theta_r)$  and  $A_r(f, \theta_r)$ ,  $r = \overline{1, N}$ . Here  $\theta_{r-1} \leq \zeta_{r-1} \leq \theta_r$  for all  $r = 1, 2, \dots, N$ .

*Step 2.* Using the found matrices and vectors compose the system of linear algebraic equations (12).

*Step 3.* Solve the constructed system and find  $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_N^*) \in \mathbb{R}^{nN}$ . Note that the elements of  $\lambda^*$  are the values of the solution to multi-point problem (1.1), (1.2) at the interior points of the subintervals:  $\lambda_r^* = u^*(\zeta_{r-1})$ ,  $r = \overline{1, N}$ .

*Step 4.* Solve the Cauchy problems

$$\frac{dz}{dt} = A(t)z + f(t), \quad z(\zeta_{r-1}) = \lambda_r^*, \quad t \in [\theta_{r-1}, \theta_r],$$

and define the values of the solution  $x^*(t)$  at the remaining points of the subintervals  $[\theta_{r-1}, \theta_r]$ ,  $r = \overline{1, N}$ .

The function  $x^*(t)$  is a solution to original multi-point problem (1.1), (1.2).

As it follows from Lemma 3.1, any solution to system (3.3) determines the values of the solution to problem (1.1), (1.2) at the left end-points of the subintervals  $[\theta_{r-1}, \theta_r]$ ,  $r = \overline{1, N}$ .

The accuracy of the algorithm proposed depends on the accuracy of computing the coefficients and right-hand sides of system of algebraic equations (3.3).

The Cauchy problem for a system of ordinary differential equations is the principal auxiliary problem in the offered algorithm. By choosing an approximate method for solving that problem, we obtain an approximate method for solving the multi-point boundary value problem (1.1), (1.2). The solution of the Cauchy problems by numerical methods leads to numerical algorithms for solving multi-point problem (1.1), (1.2).

**Remark 1.** In the general case, the points  $t = \theta_i$  in the multi-point condition may not coincide with the left-end points of the subintervals  $[\theta_{r-1}, \theta_r)$ ,  $r = \overline{1, N}$ . In this case, we can re-number all the points so that the points of the multi-point condition become the left-end points of the subintervals.

**Conclusion.** In the paper, we propose a new approach aimed at studying multi-point boundary value problems for systems of differential equations with a piecewise constant argument of generalized type. This method is based on a new concept of general solution of differential equations with piecewise constant argument of generalized type and Dzhumabaev's parametrization method. New general solution enables us to establish the qualitative properties of multi-point boundary value problems for systems of differential equations with a piecewise constant argument of generalized type and to develop algorithms for solving them. The algorithms are based on constructing and solving systems of linear algebraic equations in arbitrary vectors of new general solution. The results obtained can be used in a wide range of applications: problems for impulsive differential equations with a piecewise constant argument of generalized type; the theory of dynamical systems and neural networks; nonlocal problems for hyperbolic equations with a piecewise constant argument of generalized type, etc. [17], [36].

## Acknowledgments

The second author (Assanova) presented the results for a two-point boundary value problems for DEPCAG at the Mini-Symposium "Differential equations, dynamical systems and applications" (MS - ID 52) at the 8th European Congress of Mathematics, 20-26 June, 2021, Portorož, Slovenia [12].

This research is funded by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan (Grant No. AP08855726).



The authors (together with other colleagues) have received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 873071.

## References

- [1] M.U. Akhmet, *Integral manifolds of differential equations with piecewise constant argument of generalized type*. Nonlinear Analysis, 66 (2007), 367–383.
- [2] M.U. Akhmet, *On the reduction principle for differential equations with piecewise constant argument of generalized type*. J. Math. Anal. Appl., 336 (2007), 646–663.
- [3] M.U. Akhmet, *Almost periodic solutions of differential equations with piecewise constant argument of generalized type*. Nonl. Anal.: Hybrid Systems and Appl., 2 (2008), 456–467.
- [4] M.U. Akhmet, *Stability of differential equations with piecewise constant argument of generalized type*. Nonl. Anal.: Theory, Methods and Appl., 68 (2008), 794–803.
- [5] M.U. Akhmet, *Principles of discontinuous dynamical systems*. Springer, New-York, 2010.
- [6] M.U. Akhmet, *Nonlinear hybrid continuous/discrete-time models*. Atlantis Press, Paris, 2011.
- [7] M. Akhmet, *Almost periodicity, chaos, and asymptotic equivalence*. Springer, New-York, 2020.
- [8] M. Akhmet, M. Dauylbayev, A. Mirzakulova, *A singularly perturbed differential equation with piecewise constant argument of generalized type*. Turkish J. Math., 42 (2018), 1680–1685.
- [9] M. Akhmet, M.O. Fen, *Replication of chaos in neural networks, economics and physics. Nonlinear physical science*. Springer, Higher Education Press, Beijing, Heidelberg, 2016.
- [10] M. Akhmet, A. Kashkynbayev, *Bifurcation in autonomous and nonautonomous differential equations with discontinuities*. Springer, Higher Education Press, Heidelberg, 2017.
- [11] M.U. Akhmet, E. Yilmaz, *Neural networks with discontinuous/impact activations*. Springer, New-York, 2013.
- [12] A.T. Assanova, *A boundary value problem for system of differential equations with piecewise-constant argument of generalized type*. 8th European Congress of Mathematics. 20–26 June 2021, Portorož, Slovenia. Book of Abstracts. 152–153.
- [13] A.T. Assanova, E.A. Bakirova, Z.M. Kadirbayeva & R.E. Uteshova, *A computational method for solving a problem with parameter for linear systems of integro-differential equations*. Comput. Appl. Math., 39 (2020), Art. No. 248.
- [14] A.T. Assanova, N.B. Iskakova, N.T. Orumbayeva, *On the well-posedness of periodic problems for the system of hyperbolic equations with finite time delay*. Math. Meth. Appl. Sci., 43 (2020), 881–902.
- [15] A.T. Assanova, Z.M. Kadirbayeva, *On the numerical algorithms of parametrization method for solving a two-point boundary-value problem for impulsive systems of loaded differential equations*. Comput. Appl. Math., 37 (2018), 4966–4976.
- [16] A.T. Assanova, Z.M. Kadirbayeva, *Periodic problem for an impulsive system of the loaded hyperbolic equations*. Electr. J. Differ. Equ., 2018 (2018), no. 72, 1–8.
- [17] A.T. Assanova, Z.S. Tokmurzin, *A nonlocal multipoint problem for a system of fourth-order partial differential equations*. Eurasian Math. J., 11 (2020), no. 3, 3–20.
- [18] G. Bao, S. Wen, Z. Zeng, *Robust stability analysis of interval fuzzy Cohen–Grossberg neural networks with piecewise constant argument of generalized type*. Neural Networks, 33 (2012), 32–41.
- [19] S. Busenberg, K.L. Cooke, *Models of vertically transmitted diseases with sequential-continuous dynamics*. In "Nonlinear Phenomena in Mathematical Sciences" (V. Lakshmikantham, Ed.), Academic Press, NY, (1982), 179–187.
- [20] S. Busenberg, K.L. Cooke, *Vertically transmitted diseases. Models and dynamics*. Springer-Verlag, Berlin - Heidelberg, 1993.
- [21] S. Castillo, M. Pinto, R. Torres, *Asymptotic formulae for solutions to impulsive differential equations with piecewise constant argument of generalized type*. Electronic J. Differ. Equ., 2019 (2019), no. 40, 1–22.

- [22] K.L. Cooke, J. Wiener, *Retarded differential equations with piecewise constant delays*. J. Math. Anal. Appl., 99 (1984), 265–297.
- [23] O. Diekmann, S.A. van Gils, S.M.V. Lüne, H.-O. Walther, *Delay equations. functional-, complex-, and nonlinear analysis*. Springer-Verlag, New York, 1995.
- [24] M.A. Dominguez-Perez, R. Rodriguez-Lopez, *Multipoint BVPs of Neumann type for functional differential equations*. Nonl. Anal.: Real World Appl., 13 (2012), 1662–1675.
- [25] D.S. Dzhumabayev, *Criteria for the unique solvability of a linear boundary-value problem for an ordinary differential equation*. U.S.S.R. Comput. Math. and Math. Phys., 29 (1989), 34–46.
- [26] D.S. Dzhumabaev, *On one approach to solve the linear boundary value problems for Fredholm integro-differential equations*. J. Comput. Appl. Math., 294 (2016), 342–357.
- [27] D.S. Dzhumabaev, *New general solutions to linear Fredholm integro-differential equations and their applications on solving the BVPs*. J. Comput. Appl. Math., 327 (2018), 79–108.
- [28] D.S. Dzhumabaev, *Computational methods of solving the BVPs for the loaded differential and Fredholm integro-differential equations*. Math. Meth. Appl. Sci., 41 (2018), 1439–1462.
- [29] D.S. Dzhumabaev, *Well-posedness of nonlocal boundary-value problem for a system of loaded hyperbolic equations and an algorithm for finding its solution*. J. Math. Anal. Appl., 461 (2018), 817–836.
- [30] D.S. Dzhumabaev, *New general solutions of ordinary differential equations and the methods for the solution of boundary - value problems*. Ukrainian Math. J., 71 (2019), 1006–1031.
- [31] D.S. Dzhumabaev, E.A. Bakirova, S.T. Mynbayeva, *A method of solving a nonlinear boundary value problem with a parameter for a loaded differential equation*. Math. Meth. Appl. Sci., 43, 1788–1802 (2020).
- [32] D.S. Dzhumabaev, S.T. Mynbayeva, *New general solution to a nonlinear Fredholm integro-differential equation*. Eurasian Math. J., 10 (2019), no. 4, 24-33.
- [33] I. Gyori, G. Ladas, *Oscillation theory of delay differential equations. with applications*. Oxford Univ. Press, New York, 1991.
- [34] J. Hale, S.M.V. Lüne, *Introduction to functional differential equations*. Springer-Verlag, New York, 1993.
- [35] J.J. Nieto, R. Rodriguez-Lopez, *Monotone method for first-order functional differential equations*. Computers & Math. with Appl., 52 (2006), 471–484.
- [36] N.T. Orumbayeva, A.T. Assanova, A.B. Keldibekova, *On an algorithm of finding an approximate solution of a periodic problem for a third-order differential equation*. Eurasian Math. J., 13 (2022), no. 1, 69BТ“85.
- [37] R. Rodriguez-Lopez, *Nonlocal BVPs for second-order functional differential equations*. Nonl. Anal., 74 (2011), 7226–7239.
- [38] A.M. Samoilenko, N.A. Perestyuk, *Impulsive differential equations*. World Scientific, Singapore, 1995.
- [39] S.M. Shah, J. Wiener, *Advanced differential equations with piecewise constant argument deviations*. Intern. J. Math. and Math. Sciences, 6 (1983), 671–703.
- [40] J. Wiener, *Generalized solutions of functional differential equations*. World Scientific, Singapore, 1993.

Aziza Abildayeva, Anar Assanova, Askarbek Imanchiyev  
 Department of Mathematical Physics and Modeling  
 Institute of Mathematics and Mathematical Modeling  
 125 Pushkin St,  
 050010 Almaty, Kazakhstan  
 E-mails: azizakz@mail.ru, anartasan@gmail.com, imanchiev\_ae@mail.ru



Askarbek Imanchiyev  
Department of Mathematics  
Zhubanov Aktobe Regional University  
3 Aliya Moldagulova Ave,  
030000 Aktobe, Kazakhstan  
E-mail: imanchiev\_ae@mail.ru

Received: 09.07.2021