

ISSN 2077–9879

Eurasian Mathematical Journal

2013, Volume 4, Number 1

Founded in 2010 by
the L.N. Gumilyov Eurasian National University
in cooperation with
the M.V. Lomonosov Moscow State University
the Peoples' Friendship University of Russia
the University of Padua

Supported by the ISAAC
(International Society for Analysis, its Applications and Computation)

Published by
the L.N. Gumilyov Eurasian National University
Astana, Kazakhstan

EURASIAN MATHEMATICAL JOURNAL

Editorial Board

Editors-in-Chief

V.I. Burenkov, M. Otelbaev, V.A. Sadovnichy

Editors

Sh.A. Alimov (Uzbekistan), H. Begehr (Germany), O.V. Besov (Russia), B. Bójarski (Poland), A.A. Borubaev (Kyrgyzstan), G. Bourdaud (France), R.C. Brown (USA), A. Caetano (Portugal), M. Carro (Spain), A.D.R. Choudary (Pakistan), V.N. Chubarikov (Russia), A.S. Dzumadildaev (Kazakhstan), V.M. Filippov (Russia), H. Ghazaryan (Armenia), M.L. Goldman (Russia), V. Goldshtein (Israel), V. Guliyev (Azerbaijan), D.D. Haroske (Germany), A. Hasanoglu (Turkey), M. Huxley (Great Britain), M. Imanaliev (Kyrgyzstan), P. Jain (India), T.Sh. Kalmenov (Kazakhstan), K.K. Kenzhibaev (Kazakhstan), E. Kissin (Great Britain), V. Kokilashvili (Georgia), A. Kufner (Czech Republic), L.K. Kussainova (Kazakhstan), P.D. Lamberti (Italy), M. Lanza de Cristoforis (Italy), V.G. Maz'ya (Sweden), A.V. Mikhalev (Russia), E.D. Nursultanov (Kazakhstan), R. Oinarov (Kazakhstan), I.N. Parasidis (Greece), J. Pečarić (Croatia), S.A. Plaksa (Ukraine), L.-E. Persson (Sweden), E.L. Presman (Russia), M.D. Ramazanov (Russia), M. Reissig (Germany), S. Sagitov (Sweden), T.O. Shaposhnikova (Sweden), A.A. Shkalikov (Russia), V.A. Skvortsov (Poland), G. Sinnamon (Canada), E.S. Smailov (Kazakhstan), V.D. Stepanov (Russia), Ya.T. Sultanaev (Russia), I.A. Taimanov (Russia), T.V. Tararykova (Great Britain), U.U. Umirbaev (Kazakhstan), Z.D. Usmanov (Tajikistan), B. Viscolani (Italy), Masahiro Yamamoto (Japan), Dachun Yang (China), B.T. Zhumagulov (Kazakhstan)

Managing Editor

A.M. Temirkhanova

Executive Editor

D.T. Matin

Aims and Scope

The Eurasian Mathematical Journal (EMJ) publishes carefully selected original research papers in all areas of mathematics written by mathematicians, principally from Europe and Asia. However papers by mathematicians from other continents are also welcome.

From time to time the EMJ will also publish survey papers.

The EMJ publishes 4 issues in a year.

The language of the paper must be English only.

The contents of EMJ are indexed in Mathematical Reviews, Zentralblatt Math (ZMATH), Referativnyi Zhurnal – Matematika, Math-Net.Ru.

Information for the Authors

Submission. Manuscripts should be written in LaTeX and should be submitted electronically in DVI, PostScript or PDF format to the EMJ Editorial Office via e-mail (eurasianmj@yandex.kz).

When the paper is accepted, the authors will be asked to send the tex-file of the paper to the Editorial Office.

The author who submitted an article for publication will be considered as a corresponding author. Authors may nominate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not guaranteed.

Copyright. When the paper is accepted, the copyright is automatically transferred to the EMJ. Manuscripts are accepted for review on the understanding that the same work has not been already published (except in the form of an abstract), that it is not under consideration for publication elsewhere, and that it has been approved by all authors.

Title page. The title page should start with the title of the paper and authors' names (no degrees). It should contain the Keywords (no more than 10), the Subject Classification (AMS Mathematics Subject Classification (2010) with primary (and secondary) subject classification codes), and the Abstract (no more than 150 words with minimal use of mathematical symbols).

Figures. Figures should be prepared in a digital form which is suitable for direct reproduction.

References. Bibliographical references should be listed alphabetically at the end of the article. The authors should consult the Mathematical Reviews for the standard abbreviations of journals' names.

Authors' data. The authors' affiliations, addresses and e-mail addresses should be placed after the References.

Proofs. The authors will receive proofs only once. The late return of proofs may result in the paper being published in a later issue.

Honorarium. An honorarium will be paid to the authors (at 50 \$ per page, but no more than 2000 \$ for a paper) and to the reviewers (at 50 \$ per review). All papers will be edited with regard to usage of English. If severe editing is required, the honorarium may be reduced to 40 \$ per page. Papers may be submitted in Russian. They will be translated into English, and in this case the honorarium will be 25 \$ per page.

Offprints. The authors will receive offprints in electronic form.

Publication Ethics and Publication Malpractice

Manuscripts are accepted for review on the understanding that the same work has not been already published (except in the form of an abstract), that it is not under consideration for publication elsewhere, and that it has been approved by all authors. Only original papers are accepted for consideration. In particular, translations into English of papers already published in another language are not accepted. No other forms of scientific misconduct are allowed, such as plagiarism, falsification, incorrect interpretation of other works, incorrect citations, etc.

Web-page

The web-page of EMJ is www.emj.enu.kz. One can enter the web-page by typing Eurasian Mathematical Journal in any search engine (Google, Yandex, etc.). The archive of the web-page contains all papers published in EMJ (free access).

Subscription

For Institutions

- US\$ 200 (or equivalent) for one volume (4 issues)
- US\$ 60 (or equivalent) for one issue

For Individuals

- US\$ 160 (or equivalent) for one volume (4 issues)
- US\$ 50 (or equivalent) for one issue.

The price includes handling and postage.

The Subscription Form for subscribers can be obtained by e-mail:

eurasianmj@yandex.kz

The Eurasian Mathematical Journal (EMJ)
The Editorial Office
The L.N. Gumilyov Eurasian National University,
Main Building
Room 355
Tel.: +7-7172-709500 extension 31313
5 Munaitpasov Street
010008 Astana
Kazakhstan

Proceedings of the international workshop
OPERATORS IN MORREY-TYPE SPACES AND APPLICATIONS
(OMTSA 2011)

dedicated to 70th birthday of Professor V.I. Burenkov

held through May 20-27, 2011

at the Ahi Evran University, Kirşehir, Turkey

Volume 2

NEW CHARACTERIZATION OF MORREY SPACES

A. Gogatishvili, R.Ch. Mustafayev

Communicated by M.L. Goldman

Key words: local Morrey-type spaces, complementary local Morrey-type spaces, associate spaces, dual spaces.

AMS Mathematics Subject Classification: 26D15, 46E30.

Abstract. In this paper we prove that the norm of the Morrey space $\mathcal{M}_{p,\lambda}$ is equivalent to

$$\sup \left\{ \int_{\mathbb{R}^n} |fg| : \inf_{x \in \mathbb{R}^n} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathfrak{C}_{B(x,r)})} dr \leq 1 \right\}.$$

1 Introduction

The well-known Morrey spaces $\mathcal{M}_{p,\lambda}$ introduced by C. Morrey in 1938 [7] in relation to the study of partial differential equations, were widely investigated during last decades, including the study of classical operators of harmonic analysis - maximal, singular and potential operators - in generalizations of these spaces (so-called Morrey-type spaces). These spaces appeared to be quite useful in the study of the local behaviour of solutions to partial differential equations, a priori estimates and other topics in the theory of partial differential equations.

In [4] local Morrey-type spaces $LM_{p\theta,\omega}$ and global Morrey-type spaces $GM_{p\theta,\omega}$ were defined and some properties of these spaces were studied. Authors investigated the boundedness of the Hardy-Littlewood maximal operator in these spaces. After this paper there was intensive study of boundedness of other classical operators such as fractional maximal operator, Riesz potential and Calderón-Zygmund singular integral operator (see, for instance [2], [3], [6], for references).

Later in [5] so-called complementary local Morrey-type spaces ${}^{\mathfrak{C}}LM_{p\theta,\omega}$ were introduced and the boundedness of fractional maximal operator from a complementary local Morrey-type space ${}^{\mathfrak{C}}LM_{p\theta,\omega}$ to a local Morrey-type space $LM_{p\theta,\omega}$ was investigated. As in the definition of the space ${}^{\mathfrak{C}}LM_{p\theta,\omega}$ the complement of a ball was used instead of a ball, it was named a complementary local Morrey-type space. We note that relationship between the spaces $LM_{p\theta,\omega}$ and ${}^{\mathfrak{C}}LM_{p\theta,\omega}$ was not studied.

In [6] the associated spaces and dual spaces of local Morrey-type spaces and complementary local Morrey-type spaces were characterized. More precisely, it was shown that the associated spaces of local Morrey-type spaces are complementary local Morrey-type spaces. Moreover, it was proved that for some values of the parameters these associated spaces are duals of local Morrey-type spaces. Namely, it was proved that the space

${}^cLM_{p'\theta',\tilde{\omega}}$ is dual space of the space $LM_{p\theta,\omega}$, where $1 \leq p, \theta < \infty$, p' and θ' are the conjugate exponents of p, θ respectively, and $\tilde{\omega}(t) = \omega^{\theta-1}(t) \left(\int_t^\infty \omega^\theta(s)ds\right)^{-1}$.

In the present paper we show that similar results hold for $LM_{p\theta,\omega}^{\{x\}}$ and ${}^cLM_{p\theta,w}^{\{x\}}$ spaces for any $x \in \mathbb{R}^n$ (see Section 2 for definitions). By means of these results we prove Hölder's inequality for the classical Morrey spaces.

The paper is organized as follows. We start with notation and give some preliminaries in Section 2. In Section 3 we recall some results on associate spaces of local Morrey-type spaces and complementary local Morrey-type spaces. New characterization of the Morrey space is given in Section 4.

2 Notation and Preliminaries

Now we make some conventions. Throughout the paper, we always denote by c and C a positive constant which is independent of main parameters, but it may vary from line to line. By $A \lesssim B$ we mean that $A \leq cB$ with some positive constant c independent of the appropriate quantities. If $A \lesssim B$ and $B \lesssim A$, we write $A \sim B$ and say that A and B are equivalent. A constant with subscript, such as c_1 , does not change in different occurrences. For a measurable set E , χ_E denotes the characteristic function of E .

Unless a special remarks is made, the differential element dx is omitted when the integrals under consideration are the Lebesgue integrals.

For a fixed p with $p \in [1, \infty)$, p' denotes the conjugate exponent of p , namely,

$$p' := \begin{cases} \frac{p}{1-p} & \text{if } 0 < p < 1, \\ +\infty & \text{if } p = 1, \\ \frac{p}{p-1} & \text{if } 1 < p < +\infty, \\ 1 & \text{if } p = +\infty, \end{cases}$$

and $1/(+\infty) = 0$, $0/0 = 0$, $0 \cdot (\pm\infty) = 0$.

If E is a nonempty measurable subset on \mathbb{R}^n and f is a measurable function on E , then we put

$$\|g\|_{L_p(E)} := \left(\int_E |f(y)|^p dy\right)^{\frac{1}{p}}, \quad 0 < p < +\infty,$$

$$\|f\|_{L_\infty(E)} := \sup\{\alpha : |\{y \in E : |f(y)| \geq \alpha\}| > 0\}.$$

If I is a nonempty measurable subset on $(0, +\infty)$ and g is a measurable function on I , then we define $\|g\|_{L_p(I)}$ and $\|g\|_{L_\infty(I)}$, correspondingly.

For $x \in \mathbb{R}^n$ and $r > 0$, let $B(x, r)$ be the open ball centered at x of radius r and ${}^cB(x, r) := \mathbb{R}^n \setminus B(x, r)$.

Morrey spaces $\mathcal{M}_{p,\lambda}$ were introduced by C. Morrey in 1938 [7] and defined as follows: for $0 \leq \lambda \leq n$, $1 \leq p \leq \infty$, $f \in \mathcal{M}_{p,\lambda}$ if $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ and

$$\|f\|_{\mathcal{M}_{p,\lambda}} \equiv \|f\|_{\mathcal{M}_{p,\lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} r^{\frac{\lambda-n}{p}} \|f\|_{L_p(B(x,r))} < \infty.$$

Note that $\mathcal{M}_{p,0} = L_\infty(\mathbb{R}^n)$ and $\mathcal{M}_{p,n} = L_p(\mathbb{R}^n)$. If $\lambda < 0$ or $\lambda > n$, then $\mathcal{M}_{p,\lambda} = \Theta$, where Θ is the set of all functions equivalent to 0 on \mathbb{R}^n .

In [1] D.R.Adams introduced a variant of Morrey-type spaces as follows: For $0 \leq \lambda \leq n$, $1 \leq p, \theta \leq \infty$, $f \in \mathcal{M}_{p\theta,\lambda}$ if $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ and

$$\|f\|_{\mathcal{M}_{p\theta,\lambda}} \equiv \|f\|_{\mathcal{M}_{p\theta,\lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n} \|r^{\frac{\lambda-n}{p}} \|f\|_{L_p(B(x,r))}\|_{L_\theta(0,\infty)} < \infty.$$

(If $\theta = \infty$, then $\mathcal{M}_{p\theta,\lambda} = \mathcal{M}_{p,\lambda}$.)

Let us recall definitions of local Morrey-type space and complementary local Morrey-type space.

Definition 1. ([4]) Let $0 < p, \theta \leq \infty$ and let w be a non-negative measurable function on $(0, \infty)$. We denote by $LM_{p\theta,\omega}$ the local Morrey-type space, the space of all functions $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ with finite quasinorm

$$\|f\|_{LM_{p\theta,\omega}} \equiv \|f\|_{LM_{p\theta,\omega}(\mathbb{R}^n)} = \|w(r)\|f\|_{L_p(B(0,r))}\|_{L_\theta(0,\infty)}.$$

Definition 2. ([5]) Let $0 < p, \theta \leq \infty$ and let w be a non-negative measurable function on $(0, \infty)$. We denote by ${}^cLM_{p\theta,\omega}$ the complementary local Morrey-type space, the space of all functions $f \in L_p({}^cB(0,t))$ for all $t > 0$ with finite quasinorm

$$\|f\|_{{}^cLM_{p\theta,\omega}} \equiv \|f\|_{{}^cLM_{p\theta,\omega}(\mathbb{R}^n)} = \left\| w(r)\|f\|_{L_p({}^cB(0,r))} \right\|_{L_\theta(0,\infty)}.$$

Definition 3. Let $0 < p, \theta \leq \infty$. We denote by Ω_θ the set all non-negative measurable functions ω on $(0, \infty)$ such that

$$0 < \|\omega\|_{L_\theta(t,\infty)} < \infty, \quad t > 0,$$

and by ${}^c\Omega_\theta$ the set all non-negative measurable functions ω on $(0, \infty)$ such that

$$0 < \|\omega\|_{L_\theta(0,t)} < \infty, \quad t > 0.$$

It is convenient to define local Morrey-type spaces and complementary local Morrey-type spaces at any fixed point $x \in \mathbb{R}^n$.

Definition 4. Let $0 < p, \theta \leq \infty$ and let w be a non-negative measurable function on $(0, \infty)$. For any fixed $x \in \mathbb{R}^n$ we denote by $LM_{p\theta,\omega}^{\{x\}}$, the local Morrey-type space: the space of all functions $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ with finite quasinorm

$$\|f\|_{LM_{p\theta,\omega}^{\{x\}}} \equiv \|f\|_{LM_{p\theta,\omega}^{\{x\}}(\mathbb{R}^n)} := \|w(r)\|f\|_{L_p(B(x,r))}\|_{L_\theta(0,\infty)} = \|f(x + \cdot)\|_{LM_{p\theta,\omega}}.$$

Definition 5. Let $0 < p, \theta \leq \infty$ and let w be a non-negative measurable function on $(0, \infty)$. For any fixed $x \in \mathbb{R}^n$ we denote by ${}^cLM_{p\theta,\omega}^{\{x\}}$ the complementary local Morrey-type space, the space of all functions $f \in L_p({}^cB(x,t))$ for all $t > 0$ with finite quasinorm

$$\|f\|_{{}^cLM_{p\theta,\omega}^{\{x\}}} \equiv \|f\|_{{}^cLM_{p\theta,\omega}^{\{x\}}(\mathbb{R}^n)} := \left\| w(r)\|f\|_{L_p({}^cB(x,r))} \right\|_{L_\theta(0,\infty)} = \|f(x + \cdot)\|_{{}^cLM_{p\theta,\omega}}.$$

Clearly $LM_{p\theta,\omega} = LM_{p\theta,\omega}^{\{0\}}$ and ${}^cLM_{p\theta,\omega} = {}^cLM_{p\theta,\omega}^{\{0\}}$.

3 Associate and dual spaces of local Morrey-type and complementary local Morrey-type spaces

Let (\mathcal{R}, μ) be a totally σ -finite non-atomic measure space. Let $\mathfrak{M}(\mathcal{R}, \mu)$ be the set of all μ -measurable a.e. finite real functions on \mathcal{R} .

Definition 6. Let X be a set of functions from $\mathfrak{M}(\mathcal{R}, \mu)$, endowed with a positively homogeneous functional $\|\cdot\|_X$, defined for every $f \in \mathfrak{M}(\mathcal{R}, \mu)$ and such that $f \in X$ and only if $\|f\|_X < \infty$. We define the associate space X' of X as the set of all functions $f \in \mathfrak{M}(\mathcal{R}, \mu)$ such that $\|f\|_{X'} < \infty$, where

$$\|f\|_{X'} = \sup \left\{ \int_{\mathcal{R}} |fg| d\mu : \|g\|_X \leq 1 \right\}.$$

In what follows we assume $\mathcal{R} = \mathbb{R}^n$ and $d\mu = dx$.

In [6] the associate spaces of local Morrey-type and complementary local Morrey-type spaces were described. Our method of characterization of the Morrey space is mainly based on these results. For the sake of completeness we recall some statements from [6].

Theorem 3.1. ([6], Theorem 4.2) *Assume $1 \leq p < \infty$, $0 < \theta \leq \infty$. Let $\omega \in {}^{\circ} \Omega_{\theta}$. Set $X = {}^{\circ} LM_{p\theta, \omega}$.*

(i) Let $0 < \theta \leq 1$. Then

$$\|f\|_{X'} \sim \sup_{t \in (0, \infty)} \|f\|_{L_{p'}(B(0,t))} \|\omega\|_{L_{\theta}(0,t)}^{-1},$$

with the positive constant in the equivalency independent of f .

(ii) Let $1 < \theta \leq \infty$. Then

$$\|f\|_{X'} \sim \left(\int_{(0, \infty)} \|f\|_{L_{p'}(B(0,t))}^{\theta'} d \left(-\|\omega\|_{L_{\theta}(0,t+)}^{-\theta'} \right) \right)^{\frac{1}{\theta'}} + \frac{\|f\|_{L_{p'}(\mathbb{R}^n)}}{\|\omega\|_{L_{\theta}(0, \infty)}},$$

with the positive constant in the equivalency independent of f , where

$$\|\omega\|_{L_{\theta}(0,t+)} := \lim_{s \rightarrow t+} \|\omega\|_{L_{\theta}(0,s)}, \quad t \in (0, \infty).$$

Theorem 3.2. ([6], Theorem 4.3) *Assume $1 \leq p < \infty$, $0 < \theta \leq \infty$. Let $\omega \in \Omega_{\theta}$. Set $X = LM_{p\theta, \omega}$.*

(i) Let $0 < \theta \leq 1$. Then

$$\|f\|_{X'} \sim \sup_{t \in (0, \infty)} \|f\|_{L_{p'}({}^{\circ} B(0,t))} \|\omega\|_{L_{\theta}(t, \infty)}^{-1},$$

with the positive constant in the equivalency independent of f .

(ii) Let $1 < \theta \leq \infty$. Then

$$\|f\|_{X'} \sim \left(\int_{(0, \infty)} \|f\|_{L_{p'}({}^{\circ} B(0,t))}^{\theta'} d \|\omega\|_{L_{\theta}(t-, \infty)}^{-\theta'} \right)^{\frac{1}{\theta'}} + \frac{\|f\|_{L_{p'}(\mathbb{R}^n)}}{\|\omega\|_{L_{\theta}(0, \infty)}},$$

with the positive constant in the equivalency independent of f , where

$$\|\omega\|_{L_{\theta}(t-, \infty)} := \lim_{s \rightarrow t-} \|\omega\|_{L_{\theta}(s, \infty)}, \quad t \in (0, \infty).$$

In fact more general results, which are important for our applications, are true.

Theorem 3.3. *Assume $1 \leq p < \infty$, $0 < \theta \leq \infty$. Let $\omega \in {}^c\Omega_\theta$. For any fixed $x \in \mathbb{R}^n$ set $X = {}^cLM_{p\theta,\omega}^{\{x\}}$.*

(i) Let $0 < \theta \leq 1$. Then

$$\|f\|_{X'} \sim \sup_{t \in (0, \infty)} \|f\|_{L_{p'}(B(x,t))} \|\omega\|_{L_\theta(0,t)}^{-1},$$

with the positive constant in the equivalency independent of f and x .

(ii) Let $1 < \theta \leq \infty$. Then

$$\|f\|_{X'} \sim \left(\int_{(0, \infty)} \|f\|_{L_{p'}(B(x,t))}^{\theta'} d \left(-\|\omega\|_{L_\theta(0,t+)}^{-\theta'} \right) \right)^{\frac{1}{\theta'}} + \frac{\|f\|_{L_{p'}(\mathbb{R}^n)}}{\|\omega\|_{L_\theta(0, \infty)}},$$

with the positive constant in the equivalency independent of f and x .

Proof. Let x be any fixed point in \mathbb{R}^n . Then

$$\begin{aligned} \|f\|_{X'} &= \|f\|_{({}^cLM_{p\theta,\omega}^{\{x\}})'} = \sup \left\{ \int_{\mathbb{R}^n} |f(y)g(y)| dy : \|g\|_{{}^cLM_{p\theta,\omega}^{\{x\}}} \leq 1 \right\} \\ &= \sup \left\{ \int_{\mathbb{R}^n} |f(x+y)g(x+y)| dy : \|g(x+\cdot)\|_{{}^cLM_{p\theta,\omega}^{\{x\}}} \leq 1 \right\} \\ &= \sup \left\{ \int_{\mathbb{R}^n} |f(x+y)g(y)| dy : \|g\|_{{}^cLM_{p\theta,\omega}} \leq 1 \right\} \\ &= \|f(x+\cdot)\|_{({}^cLM_{p\theta,\omega})'}. \end{aligned}$$

It remains to apply Theorem 3.1. □

Theorem 3.4. *Assume $1 \leq p < \infty$, $0 < \theta \leq \infty$. Let $\omega \in \Omega_\theta$. For any fixed $x \in \mathbb{R}^n$ set $X = LM_{p\theta,\omega}^{\{x\}}$.*

(i) Let $0 < \theta \leq 1$. Then

$$\|f\|_{X'} \sim \sup_{t \in (0, \infty)} \|f\|_{L_{p'}({}^cB(x,t))} \|\omega\|_{L_\theta(t, \infty)}^{-1},$$

with the positive constant in the equivalency independent of f and x .

(ii) Let $1 < \theta \leq \infty$. Then

$$\|f\|_{X'} \sim \left(\int_{(0, \infty)} \|f\|_{L_{p'}({}^cB(x,t))}^{\theta'} d \|\omega\|_{L_\theta(t, \infty)}^{-\theta'} \right)^{\frac{1}{\theta'}} + \frac{\|f\|_{L_{p'}(\mathbb{R}^n)}}{\|\omega\|_{L_\theta(0, \infty)}},$$

with the positive constant in the equivalency independent of f and x .

The proof of Theorem 3.4 is similar to that of Theorem 3.3 (we only need to apply Theorem 3.2 instead of Theorem 3.1) and we omit it.

It was shown in [6] that for some values of parameters the dual spaces coincide with the associated spaces. Namely, the following theorems were proved.

Theorem 3.5. ([6], Theorem 6.1) *Assume $1 \leq p < \infty$, $1 \leq \theta < \infty$. Let $\omega \in \Omega_\theta$ and $\|\omega\|_{L_\theta(0,\infty)} = \infty$. Then*

$$(LM_{p\theta,\omega})^* = {}^c LM_{p'\theta',\tilde{\omega}}, \quad (3.1)$$

where $\tilde{\omega}(t) = \omega^{\theta-1}(t) \left(\int_t^\infty \omega^\theta(s) ds \right)^{-1}$, under the following pairing:

$$\langle f, g \rangle = \int_{\mathbb{R}^n} fg.$$

Moreover $\|f\|_{{}^c LM_{p'\theta',\tilde{\omega}}} = \sup_g \left| \int_{\mathbb{R}^n} fg \right|$, where the supremum is taken over all functions $g \in LM_{p\theta,\omega}$ with $\|g\|_{LM_{p\theta,\omega}} \leq 1$.

Theorem 3.6. ([6], Theorem 6.2) *Assume $1 \leq p < \infty$, $1 \leq \theta < \infty$. Let $\omega \in {}^c \Omega_\theta$ and $\|\omega\|_{L_\theta(0,\infty)} = \infty$. Then*

$$\left({}^c LM_{p\theta,\omega} \right)^* = LM_{p'\theta',\bar{\omega}}, \quad (3.2)$$

where $\bar{\omega}(t) = \omega^{\theta-1}(t) \left(\int_0^t \omega^\theta(s) ds \right)^{-1}$, under the following pairing:

$$\langle f, g \rangle = \int_{\mathbb{R}^n} fg.$$

Moreover $\|f\|_{LM_{p'\theta',\bar{\omega}}} = \sup_g \left| \int_{\mathbb{R}^n} fg \right|$, where the supremum is taken over all functions $g \in {}^c LM_{p\theta,\omega} : \|g\|_{{}^c LM_{p\theta,\omega}} \leq 1$.

In fact more general results hold true.

Theorem 3.7. *Assume $1 \leq p < \infty$, $1 \leq \theta < \infty$. Let $\omega \in \Omega_\theta$ and $\|\omega\|_{L_\theta(0,\infty)} = \infty$. Then for any $x \in \mathbb{R}^n$*

$$\left(LM_{p\theta,\omega}^{\{x\}} \right)^* = {}^c LM_{p'\theta',\tilde{\omega}}^{\{x\}}, \quad (3.3)$$

where $\tilde{\omega}(t) = \omega^{\theta-1}(t) \left(\int_t^\infty \omega^\theta(s) ds \right)^{-1}$, under the following pairing:

$$\langle f, g \rangle = \int_{\mathbb{R}^n} fg.$$

Moreover $\|f\|_{{}^c LM_{p'\theta',\tilde{\omega}}^{\{x\}}} = \sup_g \left| \int_{\mathbb{R}^n} fg \right|$, where the supremum is taken over all functions $g \in LM_{p\theta,\omega}^{\{x\}}$ with $\|g\|_{LM_{p\theta,\omega}^{\{x\}}} \leq 1$.

Theorem 3.8. *Assume $1 \leq p < \infty$, $1 \leq \theta < \infty$. Let $\omega \in {}^c \Omega_\theta$ and $\|\omega\|_{L_\theta(0,\infty)} = \infty$. Then for any $x \in \mathbb{R}^n$*

$$\left({}^c LM_{p\theta,\omega}^{\{x\}} \right)^* = LM_{p'\theta',\bar{\omega}}^{\{x\}}, \quad (3.4)$$

where $\bar{\omega}(t) = \omega^{\theta-1}(t) \left(\int_0^t \omega^\theta(s) ds \right)^{-1}$, under the following pairing:

$$\langle f, g \rangle = \int_{\mathbb{R}^n} fg.$$

Moreover $\|f\|_{LM_{p'\theta',\bar{\omega}}^{\{x\}}} = \sup_g \left| \int_{\mathbb{R}^n} fg \right|$, where the supremum is taken over all functions $g \in {}^c LM_{p\theta,\omega}^{\{x\}} : \|g\|_{{}^c LM_{p\theta,\omega}^{\{x\}}} \leq 1$.

The proofs of Theorem 3.7 and Theorem 3.8 are analogous to the proofs of Theorem 3.5, Theorem 3.6 respectively, and we omit them.

4 New characterization of Morrey space

In this section we give a new characterization of the classical Morrey space.

Note that

$$g \mapsto \inf_{x \in \mathbb{R}^n} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathfrak{C}_{B(x,r)})} dr$$

is a positively homogeneous functional on $\bigcup_{x \in \mathbb{R}^n} {}^{\mathfrak{C}}LM_{p', \frac{n-\lambda}{p}-1}^{\{x\}}$.

Denote by

$$\widetilde{\mathcal{M}}_{p,\lambda} := \left\{ f \in \mathfrak{M}(\mathbb{R}^n, dx) : \|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}} < \infty \right\} \quad (4.1)$$

the associate space of the set of functions $\bigcup_{x \in \mathbb{R}^n} {}^{\mathfrak{C}}LM_{p', \frac{n-\lambda}{p}-1}^{\{x\}}$, where

$$\|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}} := \sup \left\{ \int_{\mathbb{R}^n} |fg| : \inf_{x \in \mathbb{R}^n} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathfrak{C}_{B(x,r)})} dr \leq 1 \right\}. \quad (4.2)$$

To study properties of the space $\widetilde{\mathcal{M}}_{p,\lambda}$ the following Hölder's inequality for the classical Morrey spaces are useful.

Lemma 4.1. *Let $1 \leq p < \infty$ and $0 < \lambda < n$. Then the inequality*

$$\int_{\mathbb{R}^n} |fg| \leq C \|f\|_{\mathcal{M}_{p,\lambda}} \inf_{x \in \mathbb{R}^n} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathfrak{C}_{B(x,r)})} dr, \quad (4.3)$$

holds with a positive constant C independent of functions $f, g \in \mathfrak{M}(\mathbb{R}^n, dx)$.

Proof. For $\theta = \infty$ and $w(t) = t^{\frac{\lambda-n}{p}}$ Theorem 3.4 (part (ii)) implies the following inequality

$$\int_{\mathbb{R}^n} |fg| \leq C \left(\sup_{t>0} t^{\frac{\lambda-n}{p}} \|f\|_{L_p(B(x,t))} \right) \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathfrak{C}_{B(x,r)})} dr, \quad (4.4)$$

with a constant C independent of f, g and $x \in \mathbb{R}^n$. Therefore

$$\begin{aligned} \int_{\mathbb{R}^n} |fg| &\leq C \left(\sup_{x \in \mathbb{R}^n, t>0} t^{\frac{\lambda-n}{p}} \|f\|_{L_p(B(x,t))} \right) \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathfrak{C}_{B(x,r)})} dr \\ &= C \|f\|_{\mathcal{M}_{p,\lambda}} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathfrak{C}_{B(x,r)})} dr. \end{aligned} \quad (4.5)$$

In view of arbitrariness of x we arrive at (4.3). \square

Lemma 4.2. *Let $1 \leq p < \infty$ and $0 < \lambda < n$. Then*

$$\inf_{x \in \mathbb{R}^n} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathfrak{C}_{B(x,r)})} dr = 0 \quad (4.6)$$

if and only if $g = 0$ a.e. on \mathbb{R}^n .

Proof. Obviously, $\inf_{x \in \mathbb{R}^n} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathring{c}_{B(x,r)})} dr = 0$, when $g = 0$ a.e. on \mathbb{R}^n .

Now assume that $\inf_{x \in \mathbb{R}^n} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|g\|_{L_{p'}(\mathring{c}_{B(x,r)})} dr = 0$. For any fixed $R > 0$ consider the function $f = \chi_{B(0,R)}$. Obviously, $f \in \mathcal{M}_{p,\lambda}$, since $\|\chi_{B(0,R)}\|_{\mathcal{M}_{p,\lambda}} \approx R^{\lambda/p}$. Then by inequality (4.3), we have $\int_{B(0,R)} |g| = 0$, therefore, $g = 0$ a.e. on $B(0,R)$. By arbitrariness of R , we get that $g \in \Theta$. \square

Lemma 4.3. *Let $1 \leq p < \infty$ and $0 < \lambda < n$. Then*

$$\bigcup_{x \in \mathbb{R}^n} \mathring{c}LM_{p', \frac{n-\lambda}{p}-1}^{\{x\}} \subset L_1^{\text{loc}}(\mathbb{R}^n).$$

Proof. Let g be any function in $\bigcup_{x \in \mathbb{R}^n} \mathring{c}LM_{p', \frac{n-\lambda}{p}-1}^{\{x\}}$. Then there exists $x \in \mathbb{R}^n$ such that $g \in \mathring{c}LM_{p', \frac{n-\lambda}{p}-1}^{\{x\}}$. Let R be any fixed positive number. Since the function $f = \chi_{B(x,R)} \in \mathcal{M}_{p,\lambda}$ and $\|f\|_{\mathcal{M}_{p,\lambda}} \approx R^{\lambda/p}$, by the inequality (4.5) we get

$$\int_{B(x,R)} |g(y)| dy \leq CR^{\frac{\lambda}{p}} \|g\|_{\mathring{c}LM_{p', \frac{n-\lambda}{p}-1}^{\{x\}}} < \infty.$$

In view of arbitrariness of R we get that $g \in L_1^{\text{loc}}(\mathbb{R}^n)$. \square

Lemma 4.4. *Assume $1 \leq p < \infty$ and $0 < \lambda < n$. Moreover, let $f \in L_{p'}^{\text{loc}}(\mathbb{R}^n)$. Then for any fixed $x \in \mathbb{R}^n$ and $R > 0$*

$$f\chi_{B(x,R)} \in \mathring{c}LM_{p', \frac{n-\lambda}{p}-1}^{\{x\}}.$$

Proof. Indeed, for any fixed $x \in \mathbb{R}^n$ and $R : 0 < R < \infty$, we get

$$\begin{aligned} \|f\chi_{B(x,R)}\|_{\mathring{c}LM_{p', \frac{n-\lambda}{p}-1}^{\{x\}}} &= \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|f\chi_{B(x,R)}\|_{L_{p'}(\mathring{c}_{B(x,r)})} dr \\ &= \int_0^\infty r^{\frac{n-\lambda}{p}-1} \left(\int_{\mathring{c}_{B(x,r)} \cap B(x,R)} |f|^{p'} \right)^{\frac{1}{p'}} dr \\ &= \int_0^R r^{\frac{n-\lambda}{p}-1} \left(\int_{\mathring{c}_{B(x,r)} \cap B(x,R)} |f|^{p'} \right)^{\frac{1}{p'}} dr \\ &\leq \left(\int_{B(x,R)} |f|^{p'} \right)^{\frac{1}{p'}} \int_0^R r^{\frac{n-\lambda}{p}-1} dr \\ &= c_1 R^{\frac{n-\lambda}{p}} \left(\int_{B(x,R)} |f|^{p'} \right)^{\frac{1}{p'}} < \infty. \end{aligned}$$

\square

Our main result in this section reads as follows.

Theorem 4.1. *Assume $1 \leq p < \infty$ and $0 < \lambda < n$. Then*

$$\|f\|_{\mathcal{M}_{p,\lambda}} \approx \|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}}. \quad (4.7)$$

Proof. By Lemma 4.1, it is easy to see that

$$\|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}} \lesssim \|f\|_{\mathcal{M}_{p,\lambda}}.$$

Let us prove the opposite estimate $\|f\|_{\mathcal{M}_{p,\lambda}} \lesssim \|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}}$. If $\|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}} = \infty$, then there is nothing to prove. Assume that $\|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}} < \infty$.

Observe that for $g \in L_{p'}^{\text{loc}}(\mathbb{R}^n)$ the inequality

$$\int_{B(x,R)} |fg| \leq CR^{\frac{n-\lambda}{p}} \left(\int_{B(x,R)} |g|^{p'} \right)^{\frac{1}{p'}} \|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}} \quad (4.8)$$

holds with a constant $C > 0$ independent of f , g , x and R . Indeed, let x be any fixed point in \mathbb{R}^n and $R > 0$. When $\int_{B(x,R)} |g|^{p'} = 0$ there is nothing to prove, since in this case $g = 0$ a.e. on $B(x, R)$. Assume that $\int_{B(x,R)} |g|^{p'} > 0$. Denote

$$h(y) = \frac{g(y)\chi_{B(x,R)}(y)}{c_1 R^{\frac{n-\lambda}{p}} \left(\int_{B(x,R)} |g|^{p'} \right)^{\frac{1}{p'}}}. \quad (4.9)$$

By Lemma 4.4

$$h \in {}^cLM_{p', \frac{n-\lambda}{p}-1}^{\{x\}},$$

and moreover, $\|h\|_{{}^cLM_{p', \frac{n-\lambda}{p}-1}^{\{x\}}} \leq 1$. Consequently,

$$\inf_{x \in \mathbb{R}^n} \int_0^\infty r^{\frac{n-\lambda}{p}-1} \|h\|_{L_{p'}({}^cB(x,r))} dr \leq 1.$$

Therefore

$$\int_{\mathbb{R}^n} |hf| \leq \|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}}, \quad (4.10)$$

and from (4.9), we get (4.8).

Inequality (4.8) implies that $f \in L_1^{\text{loc}}(\mathbb{R}^n)$. By Theorem of Resonance (see [8, Lemma 27, p. 283]) we get that $f \in L_p^{\text{loc}}(\mathbb{R}^n)$. The function $g := |f|^{p-1}\chi_{B(x,R)} \in L_{p'}^{\text{loc}}(\mathbb{R}^n)$, and if we put the function g in inequality (4.8), we obtain

$$\int_{B(x,R)} |f|^p \leq cR^{\frac{n-\lambda}{p}} \left(\int_{B(x,R)} |f|^p \right)^{\frac{1}{p'}} \|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}}.$$

Therefore,

$$R^{\frac{\lambda-n}{p}} \left(\int_{B(x,R)} |f|^p \right)^{\frac{1}{p}} \leq c \|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}}.$$

Since a constant c is independent of x and R , we get

$$\|f\|_{\mathcal{M}_{p,\lambda}} \leq c \|f\|_{\widetilde{\mathcal{M}}_{p,\lambda}}.$$

□

Acknowledgments

The research of A. Gogatishvili was partially supported by the grants 201/08/0383 and 13-14743S of the Grant Agency of the Czech Republic and RVO: 67985840. The research of R.Ch. Mustafayev was supported by the Science Development Foundation under the President of the Republic of Azerbaijan Project No. EIF-2010-1(1)-40/06-1.

We thank the anonymous referee for his/her remarks, which have improved the final version of this paper.

References

- [1] D.R. Adams, *Lectures on L^p -Potential Theory*, Umea Univ. Report 2 (1981), 1–74
- [2] V.I. Burenkov, *Recent progress in studying the boundedness of classical operators of real analysis in general Morrey-type spaces. I*, Eurasian Math. J. 3 (2012), no. 3, 11–32.
- [3] V.I. Burenkov, *Recent progress in studying the boundedness of classical operators of real analysis in general Morrey-type spaces. II*, Eurasian Math. J. 4 (2013), no. 1, 21–45.
- [4] V.I. Burenkov, H.V. Guliyev, *Necessary and sufficient conditions for boundedness of the maximal operator in local Morrey-type spaces*, Studia Math. 163 (2004), no. 2, 157–176.
- [5] V.I. Burenkov, H.V. Guliyev, V.S. Guliyev, *On boundedness of the fractional maximal operator from complementary Morrey-type spaces to Morrey-type spaces*, The interaction of analysis and geometry, Contemp. Math. Vol. 424, Amer. Math. Soc., Providence, RI, 2007, 17–32.
- [6] A. Gogatishvili, R.Ch. Mustafayev, *Dual spaces of local Morrey-type spaces*, Czechoslovak Math. J. 61(136) (2011), no. 3, 609–622.
- [7] C.B. Morrey, Jr. *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. 43 (1938), no. 1, 126–166.
- [8] H.L. Royden, *Real analysis*, 3rd ed., Macmillan Publishing Company, New York, 1988.

Amiran Gogatishvili

Institute of Mathematics of the Academy of Sciences of the Czech Republic

Žitna 25, 115 67 Prague 1, Czech Republic

E-mail: gogatish@math.cas.cz

Rza Mustafayev

Department of Mathematics, Faculty of Science and Arts, Kirikkale University

71450 Yahsihan, Kirikkale, Turkey

and

Institute of Mathematics and Mechanics, Academy of Sciences of Azerbaijan

F. Agayev St. 9, Baku, AZ 1141, Azerbaijan

E-mail: rzamustafayev@gmail.com

Received: 03.10.2011