

**UNIQUENESS OF THE SOLUTION TO INVERSE SCATTERING  
PROBLEM WITH BACKSCATTERING DATA**

**A.G. Ramm**

Communicated by M. Otelbaev

**Key words:** inverse scattering, non-overdetermined data, backscattering.

**AMS Mathematics Subject Classification:** 35P25, 35R30, 81Q05.

**Abstract.** Let  $q(x)$  be real-valued compactly supported sufficiently smooth function. It is proved that the scattering data  $A(-\beta, \beta, k) \forall \beta \in S^2, \forall k > 0$ , determine  $q$  uniquely.

## 1 Introduction

The scattering solution  $u(x, \alpha, k)$  solves the scattering problem:

$$[\nabla^2 + k^2 - q(x)]u = 0 \quad \text{in } \mathbb{R}^3, \quad (1)$$

$$u = e^{ik\alpha \cdot x} + A(\beta, \alpha, k) \frac{e^{ikr}}{r} + o\left(\frac{1}{r}\right), \quad r := |x| \rightarrow \infty, \quad \beta := \frac{x}{r}. \quad (2)$$

Here  $\alpha, \beta \in S^2$  are the unit vectors,  $S^2$  is the unit sphere, the coefficient  $A(\beta, \alpha, k)$  is called the scattering amplitude,  $q(x)$  is a real-valued compactly supported sufficiently smooth function. The inverse scattering problem of interest is to determine  $q(x)$  given the backscattering data  $A(-\beta, \beta, k) \forall \beta \in S^2, \forall k > 0$ . This problem is called *the inverse scattering problem with backscattering data*.

The function  $A(-\beta, \beta, k)$  depends on one unit vector  $\beta$  and on the scalar  $k$ , i.e., on three variables. The potential  $q(x)$  depends also on three variables  $x \in \mathbb{R}^3$ . This inverse problem is, therefore, not over-determined in the sense that the data and the unknown  $q(x)$  are functions of the same number of variables.

*Assumption A):* We assume that  $q$  is compactly supported, i.e.,  $q(x) = 0$  for  $|x| > a$ , where  $a > 0$  is an arbitrary large fixed number;  $q(x)$  is real-valued, i.e.,  $q = \bar{q}$ ; and  $q(x) \in H_0^\ell(B_a)$ ,  $\ell > 3$ .

Here  $B_a$  is the ball centered at the origin and of radius  $a$ , and  $H_0^\ell(B_a)$  is the closure of  $C_0^\infty(B_a)$  in the norm of the Sobolev space  $H^\ell(B_a)$  of functions whose derivatives up to the order  $\ell$  belong to  $L^2(B_a)$ .

It was proved in [5] that if  $q = \bar{q}$  and  $q \in L^2(B_a)$  is compactly supported, then the resolvent kernel  $G(x, y, k)$  of the Schrödinger operator  $-\nabla^2 + q(x) - k^2$  is a meromorphic function of  $k$  on the whole complex plane  $k$ , analytic in  $\text{Im}k \geq 0$ , except, possibly, of a finitely many simple poles at the points  $ik_j$ ,  $k_j > 0$ ,  $1 \leq j \leq n$ , where  $-k_j^2$  are negative eigenvalues of the selfadjoint operator  $-\nabla^2 + q(x)$  in  $L^2(\mathbb{R}^3)$ . Consequently,

the scattering amplitude  $A(\beta, \alpha, k)$ , corresponding to the above  $q$ , is a restriction to the positive semiaxis  $k \in [0, \infty)$  of a meromorphic on the whole complex  $k$ -plane function.

It was proved by the author ([6]), that the *fixed-energy scattering data*  $A(\beta, \alpha) := A(\beta, \alpha, k_0)$ ,  $k_0 = \text{const} > 0$ ,  $\forall \beta \in S_1^2$ ,  $\forall \alpha \in S_2^2$ , determine real-valued compactly supported  $q \in L^2(B_a)$  uniquely. Here  $S_j^2$ ,  $j = 1, 2$ , are arbitrary small open subsets of  $S^2$  (solid angles).

In [9] (see also monograph [10], Chapter 5, and [7]) an analytical formula is derived for the reconstruction of the potential  $q$  from exact fixed-energy scattering data, and from noisy fixed-energy scattering data, and stability estimates and error estimates for the reconstruction method are obtained. To the author's knowledge, these are the only known until now theoretical error estimates for the recovery of the potential from noisy fixed-energy scattering data in the three-dimensional inverse scattering problem.

In [8] stability results are obtained for the inverse scattering problem for obstacles.

The scattering data  $A(\beta, \alpha)$  depend on four variables (two unit vectors), while the unknown  $q(x)$  depends on three variables. In this sense the inverse scattering problem, which consists of finding  $q$  from the fixed-energy scattering data  $A(\beta, \alpha)$ , is overdetermined.

*Historical remark.* In the beginning of the forties of the last century physicists raised the the following question: is it possible to recover the Hamiltonian of a quantum-mechanical system from the observed quantities, such as  $S$ -matrix? In the non-relativistic quantum mechanics the simplest Hamiltonian  $\mathbf{H} = -\nabla^2 + q(x)$  can be uniquely determined if one knows the potential  $q(x)$ . The  $S$ -matrix in this case is in one-to-one correspondence with the scattering amplitude  $A$ :  $S = I - \frac{k}{2\pi i} A$ , where  $I$  is the identity operator in  $L^2(S^2)$ ,  $A$  is an integral operator in  $L^2(S^2)$  with the kernel  $A(\beta, \alpha, k)$ , and  $k^2 > 0$  is energy. Therefore, the question, raised by the physicists, is reduced to an inverse scattering problem: can one determine the potential  $q(x)$  from the knowledge of the scattering amplitude. We have briefly discussed this problem above.

Since the above question was raised, there were no uniqueness theorems for three-dimensional inverse scattering problems with non-overdetermined data. The goal of this paper is to prove such a theorem.

**Theorem 1.** *If Assumption A) holds, then the data  $A(-\beta, \beta, k) \forall \beta \in S^2, \forall k > 0$ , determine  $q$  uniquely.*

**Remark.** *The conclusion of Theorem 1 remains valid if the data  $A(-\beta, \beta, k)$  are known  $\forall \beta \in S_1^2$  and  $k \in (k_0, k_1)$ , where  $(k_0, k_1) \subset [0, \infty)$  is an arbitrary small interval,  $k_1 > k_0$ , and  $S_1^2$  is an arbitrary small open subset of  $S^2$ .*

In Section 2 we formulate some known auxiliary results.

In Section 3 proof of Theorem 1 is given.

In the Appendix a technical estimate is proved.

A brief announcement of the result is given in [3]. Although we follow the outline of the ideas from [3], the current paper is essentially self-contained and contains new arguments.

## 2 Auxiliary results

Let

$$F(g) := \tilde{g}(\xi) = \int_{\mathbb{R}^3} g(x) e^{i\xi \cdot x} dx, \quad g(x) = \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} e^{-i\xi \cdot x} \tilde{g}(\xi) d\xi. \quad (3)$$

If  $f * g := \int_{\mathbb{R}^3} f(x-y)g(y)dy$ , then

$$F(f * g) = \tilde{f}(\xi)\tilde{g}(\xi), \quad F(f(x)g(x)) = \frac{1}{(2\pi)^3} \tilde{f} * \tilde{g}. \quad (4)$$

If

$$G(x-y, k) := \frac{e^{ik[|x-y| - \beta \cdot (x-y)]}}{4\pi|x-y|}, \quad (5)$$

then

$$F(G(x, k)) = \frac{1}{\xi^2 - 2k\beta \cdot \xi}, \quad \xi^2 := \xi \cdot \xi. \quad (6)$$

The scattering solution  $u = u(x, \alpha, k)$  solves (uniquely) the integral equation

$$u(x, \alpha, k) = e^{ik\alpha \cdot x} - \int_{B_a} g(x, y, k)q(y)u(y, \alpha, k)dy, \quad (7)$$

where

$$g(x, y, k) := \frac{e^{ik|x-y|}}{4\pi|x-y|}. \quad (8)$$

If

$$v = e^{-ik\alpha \cdot x}u(x, \alpha, k), \quad (9)$$

then

$$v = 1 - \int_{B_a} G(x-y, k)q(y)v(y, \alpha, k)dy, \quad (10)$$

where  $G$  is defined in (5).

Define  $\epsilon$  by the formula

$$v = 1 + \epsilon. \quad (11)$$

Then (10) can be rewritten as

$$\epsilon(x, \alpha, k) = - \int_{\mathbb{R}^3} G(x-y, k)q(y)dy - T\epsilon, \quad (12)$$

where

$$T\epsilon := \int_{B_a} G(x-y, k)q(y)\epsilon(y, \alpha, k)dy.$$

Fourier transform of (12) yields (see (4),(6)):

$$\tilde{\epsilon}(\xi, \alpha, k) = - \frac{\tilde{q}(\xi)}{\xi^2 - 2k\alpha \cdot \xi} - \frac{1}{(2\pi)^3} \frac{1}{\xi^2 - 2k\alpha \cdot \xi} \tilde{q} * \tilde{\epsilon}. \quad (13)$$

An essential ingredient of our proof in Section 3 is the following lemma, proved by the author in [10], p.262, and in [9]. For convenience of the reader a short proof of this lemma is given in Appendix.

**Lemma 1.** *If  $A_j(\beta, \alpha, k)$  is the scattering amplitude corresponding to potential  $q_j$ ,  $j = 1, 2$ , then*

$$-4\pi[A_1(\beta, \alpha, k) - A_2(\beta, \alpha, k)] = \int_{B_1} [q_1(x) - q_2(x)]u_1(x, \alpha, k)u_2(x, -\beta, k)dx, \quad (14)$$

where  $u_j$  is the scattering solution corresponding to  $q_j$ .

Consider an algebraic variety  $\mathcal{M}$  in  $\mathbb{C}^3$  defined by the equation

$$\mathcal{M} := \{\theta \cdot \theta = 1, \quad \theta \cdot \theta := \theta_1^2 + \theta_2^2 + \theta_3^2, \quad \theta_j \in \mathbb{C}, \quad 1 \leq j \leq 3.\} \quad (15)$$

This is a non-compact variety, intersecting  $\mathbb{R}^3$  over the unit sphere  $S^2$ .

Let  $\mathbb{R}_+ = [0, \infty)$ . The following result is proved in [11], p.62.

**Lemma 2.** *If Assumption A) holds, then the scattering amplitude  $A(\beta, \alpha, k)$  is a restriction to  $S^2 \times S^2 \times \mathbb{R}_+$  of a function  $A(\theta', \theta, k)$  on  $\mathcal{M} \times \mathcal{M} \times \mathbb{C}$ , analytic on  $\mathcal{M} \times \mathcal{M}$  and meromorphic on  $\mathbb{C}$ ,  $\theta', \theta \in \mathcal{M}$ ,  $k \in \mathbb{C}$ .*

The scattering solution  $u(x, \alpha, k)$  is a meromorphic function of  $k$  in  $\mathbb{C}$ , analytic in  $\text{Im}k \geq 0$ , except, possibly, at the points  $k = ik_j$ ,  $1 \leq j \leq n$ ,  $k_j > 0$ , where  $-k_j^2$  are negative eigenvalues of the selfadjoint Schrödinger operator, defined by the potential  $q$  in  $L^2(\mathbb{R}^3)$ . These eigenvalues can be absent, for example, if  $q \geq 0$ .

We need the notion of the Radon transform:

$$\hat{f}(\beta, \lambda) := \int_{\beta \cdot x = \lambda} f(x) d\sigma, \quad (16)$$

where  $d\sigma$  is the element of the area of the plane  $\beta \cdot x = \lambda$ ,  $\beta \in S^2$ ,  $\lambda$  is a real number. The following properties of the Radon transform will be used:

$$\int_{B_a} f(x) dx = \int_{-a}^a \hat{f}(\beta, \lambda) d\lambda, \quad (17)$$

$$\int_{B_a} e^{ik\beta \cdot x} f(x) dx = \int_{-a}^a e^{ik\lambda} \hat{f}(\beta, \lambda) d\lambda, \quad (18)$$

$$\hat{f}(\beta, \lambda) = \hat{f}(-\beta, -\lambda). \quad (19)$$

These properties are proved, e.g., in [12], pp. 12, 15. We also need the following Phragmen-Lindelöf lemma, which is proved in [1], p.69, and in [2].

**Lemma 3.** *Let  $f(z)$  be holomorphic inside an angle  $\mathcal{A}$  of opening  $< \pi$ ;  $|f(z)| \leq c_1 e^{c_2|z|}$ ,  $z \in \mathcal{A}$ ,  $c_1, c_2 > 0$  are constants;  $|f(z)| \leq M$  on the boundary of  $\mathcal{A}$ ; and  $f$  is continuous up to the boundary of  $\mathcal{A}$ . Then  $|f(z)| \leq M$ ,  $\forall z \in \mathcal{A}$ .*

### 3 Proof of Theorem 1

The scattering data in Remark determine uniquely the scattering data in Theorem 1 by Lemma 2.

*Let us outline the ideas of the proof of Theorem 1.*

Assume that potentials  $q_j$ ,  $j = 1, 2$ , generate the same scattering data:

$$A_1(-\beta, \beta, k) = A_2(-\beta, \beta, k) \quad \forall \beta \in S^2, \quad \forall k > 0,$$

and let

$$p(x) := q_1(x) - q_2(x).$$

Then by Lemma 1, see equation (14), one gets

$$0 = \int_{B_a} p(x) u_1(x, \beta, k) u_2(x, \beta, k) dx, \quad \forall \beta \in S^2, \quad \forall k > 0. \quad (20)$$

By (9) and (11) one can rewrite (20) as

$$\int_{B_a} e^{2ik\beta \cdot x} [1 + \epsilon(x, k)] p(x) dx = 0 \quad \forall \beta \in S^2, \quad \forall k > 0, \quad (21)$$

where

$$\epsilon(x, k) := \epsilon := \epsilon_1(x, k) + \epsilon_2(x, k) + \epsilon_1(x, k)\epsilon_2(x, k).$$

By Lemma 2 the relations (20) and (21) hold for complex  $k$ ,

$$k = \frac{\kappa + i\eta}{2}, \quad \kappa + i\eta \neq 2ik_j, \quad \eta \geq 0. \quad (22)$$

Using formulas (3)-(4), one derives from (21) the relation

$$\tilde{p}((\kappa + i\eta)\beta) + \frac{1}{(2\pi)^3} (\tilde{\epsilon} * \tilde{p})((\kappa + i\eta)\beta) = 0 \quad \forall \beta \in S^2, \quad \forall \kappa \in \mathbb{R}, \quad (23)$$

where the notation  $(f * g)(z)$  means that the convolution  $f * g$  is calculated at the argument  $z = (\kappa + i\eta)\beta$ .

One has

$$\sup_{\beta \in S^2} |\tilde{\epsilon} * \tilde{p}| := \sup_{\beta \in S^2} \left| \int_{\mathbb{R}^3} \tilde{\epsilon}((\kappa + i\eta)\beta - s) \tilde{p}(s) ds \right| \leq \nu(\kappa, \eta) \sup_{s \in \mathbb{R}^3} |\tilde{p}(s)|, \quad (24)$$

where

$$\nu(\kappa, \eta) := \sup_{\beta \in S^2} \int_{\mathbb{R}^3} |\tilde{\epsilon}((\kappa + i\eta)\beta - s)| ds.$$

We prove that if  $\eta = \eta(\kappa) = O(\ln \kappa)$  is suitably chosen, namely as in (29) below, then the following inequality holds:

$$0 < \nu(\kappa, \eta(\kappa)) < 1, \quad \kappa \rightarrow \infty. \quad (25)$$

We also prove that

$$\sup_{\beta \in S^2} |\tilde{p}((\kappa + i\eta(\kappa))\beta)| \geq \sup_{s \in \mathbb{R}^3} |\tilde{p}(s)|, \quad \kappa \rightarrow \infty, \quad (26)$$

and then it follows from (23)-(26) that  $\tilde{p}(s) = 0$ , so  $p(x) = 0$ , and Theorem 1 is proved. Indeed, it follows from (23) and (26) that, for sufficiently large  $\kappa$  and a suitable  $\eta(\kappa) = O(\ln \kappa)$ , one has

$$\sup_{s \in \mathbb{R}^3} |\tilde{p}(s)| \leq \frac{1}{(2\pi)^3} \nu(\kappa, \eta(\kappa)) \sup_{s \in \mathbb{R}^3} |\tilde{p}(s)|.$$

If (25) holds, then the above equation implies that  $\tilde{p} = 0$ . This and the injectivity of the Fourier transform imply that  $p = 0$ .

*This completes the outline of the proof of Theorem 1.*

Let us now give a detailed proof of estimates (25) and (26), that completes the proof of Theorem 1.

We assume that  $p(x) \not\equiv 0$ , because otherwise there is nothing to prove. Let

$$\max_{s \in \mathbb{R}^3} |\tilde{p}(s)| := \mathcal{P} \neq 0.$$

**Lemma 4.** *If Assumption A) holds and  $\mathcal{P} \neq 0$ , then*

$$\limsup_{\eta \rightarrow \infty} \max_{\beta \in S^2} |\tilde{p}((\kappa + i\eta)\beta)| = \infty, \quad (27)$$

where  $\kappa > 0$  is arbitrary but fixed. For any  $\kappa > 0$  there is an  $\eta = \eta(\kappa)$ , such that

$$\max_{\beta \in S^2} |\tilde{p}((\kappa + i\eta(\kappa))\beta)| = \mathcal{P}, \quad (28)$$

where the number  $\mathcal{P} := \max_{s \in \mathbb{R}^3} |\tilde{p}(s)|$ , and

$$\eta(\kappa) = a^{-1} \ln \kappa + O(1) \quad \text{as} \quad \kappa \rightarrow +\infty. \quad (29)$$

**Proof of Lemma 4.** By formula (18) one gets

$$\tilde{p}((\kappa + i\eta)\beta) = \int_{B_a} p(x) e^{i(\kappa + i\eta)\beta \cdot x} dx = \int_{-a}^a e^{i\kappa\lambda - \eta\lambda} \hat{p}(\beta, \lambda) d\lambda. \quad (30)$$

The function  $\hat{p}(\beta, \lambda)$  is compactly supported, real-valued, and satisfies relation (19). Therefore

$$\max_{\beta \in S^2} |\tilde{p}((\kappa + i\eta(\kappa))\beta)| = \max_{\beta \in S^2} |\tilde{p}((\kappa - i\eta(\kappa))\beta)|. \quad (31)$$

Indeed,

$$\begin{aligned} \max_{\beta \in S^2} |\tilde{p}((\kappa + i\eta(\kappa))\beta)| &= \max_{\beta \in S^2} \left| \int_{-a}^a e^{i\kappa\lambda - \eta\lambda} \hat{p}(\beta, \lambda) d\lambda \right| \\ &= \max_{\beta \in S^2} \left| \int_{-a}^a e^{-i\kappa\mu + \eta\mu} \hat{p}(\beta, -\mu) d\mu \right| \\ &= \max_{\beta' \in S^2} \left| \int_{-a}^a e^{-i\kappa\mu + \eta\mu} \hat{p}(-\beta', -\mu) d\mu \right| \\ &= \max_{\beta' \in S^2} \left| \int_{-a}^a e^{-i\kappa\mu + \eta\mu} \hat{p}(\beta', \mu) d\mu \right| \\ &= \max_{\beta \in S^2} |\tilde{p}((\kappa - i\eta)\beta)|. \end{aligned} \quad (32)$$

At the last step we took into account that  $\hat{p}(\beta, \lambda)$  is a real-valued function, so

$$\begin{aligned} \max_{\beta \in S^2} \left| \int_{-a}^a e^{-i\kappa\mu + \eta\mu} \hat{p}(\beta, \mu) d\mu \right| &= \max_{\beta \in S^2} \left| \int_{-a}^a e^{i\kappa\mu + \eta\mu} \hat{p}(\beta, \mu) d\mu \right| \\ &= \max_{\beta \in S^2} |\tilde{p}((\kappa - i\eta)\beta)|. \end{aligned} \quad (33)$$

If  $p(x) \not\equiv 0$ , then (30) and (31) imply (27), as follows from Lemma 3. Let us give a detailed proof of this statement.

Consider the function  $h$  of the complex variable  $z := \kappa + i\eta$ :

$$h := h(z, \beta) := \int_{-a}^a e^{iz\lambda} \hat{p}(\beta, \lambda) d\lambda. \quad (34)$$

If (27) is false, then

$$|h(z, \beta)| \leq c \quad \forall z = \kappa + i\eta, \quad \eta \geq 0, \quad \forall \beta \in S^2, \quad (35)$$

where  $\kappa \geq 0$  is an arbitrary fixed number and the constant  $c > 0$  does not depend on  $\beta$  and  $\eta$ .

Thus,  $|h|$  is bounded on the ray  $\{\kappa = 0, \eta \geq 0\}$ , which is part of the boundary of the right angle  $\mathcal{A}$ , and the other part of its boundary is the ray  $\{\kappa \geq 0, \eta = 0\}$ . Let us check that  $|h|$  is bounded on this ray also.

One has

$$|h(\kappa, \beta)| = \left| \int_{-a}^a e^{i\kappa\lambda} \hat{p}(\beta, \lambda) d\lambda \right| \leq \int_{-a}^a |\hat{p}(\beta, \lambda)| d\lambda \leq c, \quad (36)$$

where  $c$  stands in this paper for *various* constants. From (35)-(36) it follows that on the boundary of the right angle  $\mathcal{A}$ , namely, on the two rays  $\{\kappa \geq 0, \eta = 0\}$  and  $\{\kappa = 0, \eta \geq 0\}$  the entire function  $h(z, \beta)$  of the complex variable  $z$  is bounded,  $|h(z, \beta)| \leq c$ , and inside  $\mathcal{A}$  this function satisfies the estimate

$$|h(z, \beta)| \leq e^{a|\eta|} \int_{-a}^a |\hat{p}(\beta, \lambda)| d\lambda \leq ce^{a|\eta|}, \quad (37)$$

where  $c$  does not depend on  $\beta$ . Therefore, by Lemma 3,  $|h(z, \beta)| \leq c$  in the whole angle  $\mathcal{A}$ .

By (31) the same argument is applicable to the remaining three right angles, the union of which is the whole complex  $z$ -plane  $\mathbb{C}$ . Therefore

$$\sup_{z \in \mathbb{C}, \beta \in S^2} |h(z, \beta)| \leq c. \quad (38)$$

This implies by the Liouville theorem that  $h(z, \beta) = c \forall z \in \mathbb{C}$ .

Since  $\hat{p}(\beta, \lambda) \in L^1(-a, a)$ , the relation

$$\int_{-a}^a e^{iz\lambda} \hat{p}(\beta, \lambda) d\lambda = c \quad \forall z \in \mathbb{C}, \quad (39)$$

and the Riemann-Lebesgue lemma imply that  $c = 0$ , so  $\hat{p}(\beta, \lambda) = 0 \forall \beta \in S^2$  and  $\forall \lambda \in \mathbb{R}$ . Therefore  $p(x) = 0$ , contrary to our assumption. Consequently, relation (27) is proved.

Relation (28) follows from (27) because for large  $\eta$  the left-hand side of (28) is larger than  $\mathcal{P}$  due to (27), while for  $\eta = 0$  the left-hand side of (28) is not larger than  $\mathcal{P}$  by the definition of the Fourier transform.

Let us derive estimate (29).

From the assumption  $p(x) \in H_0^\ell(B_a)$  it follows that

$$|\tilde{p}((\kappa + i\eta)\beta)| \leq c \frac{e^{a|\eta|}}{(1 + \kappa^2 + \eta^2)^{\ell/2}}. \quad (40)$$

This inequality is proved in Lemma 5, below.

The right-hand side of this inequality is of the order  $O(1)$  as  $\kappa \rightarrow \infty$  if  $|\eta| = a^{-1} \ln \kappa + O(1)$  as  $\kappa \rightarrow \infty$ . This proves relation (29) and we specify  $O(\ln \kappa)$  as in this relation.

Let us now prove inequality (40).

**Lemma 5.** *If  $p \in H_0^\ell(B_a)$  then estimate (40) holds.*

**Proof.** Consider  $\partial_j p := \frac{\partial p}{\partial x_j}$ . One has

$$\left| \int_{B_a} \partial_j p e^{i(\kappa + i\eta)\beta \cdot x} dx \right| = \left| -i(\kappa + i\eta)\beta_j \int_{B_a} p(x) e^{i(\kappa + i\eta)\beta \cdot x} dx \right| \quad (41)$$

$$= (\kappa^2 + \eta^2)^{1/2} |\tilde{p}((\kappa + i\eta)\beta)|.$$

The left-hand side of the above formula admits the following estimate

$$\left| \int_{B_a} \partial_j p e^{i(\kappa + i\eta)\beta \cdot x} dx \right| \leq c e^{a|\eta|},$$

where the constant  $c > 0$  is proportional to  $\|\partial_j p\|_{L^2(B_a)}$ . Therefore,

$$|\tilde{p}((\kappa + i\eta)\beta)| \leq c [1 + (\kappa^2 + \eta^2)]^{-1/2} e^{a|\eta|}. \quad (42)$$

Repeating this argument one gets estimate (40). Lemma 5 is proved.  $\square$

Estimate (42) implies that if relation (29) holds and  $\kappa \rightarrow \infty$ , then the quantity  $\sup_{\beta \in S^2} |\tilde{p}((\kappa + i\eta)\beta)|$  remains bounded as  $\kappa \rightarrow \infty$ .

If  $\eta$  is fixed and  $\kappa \rightarrow \infty$ , then  $\sup_{\beta \in S^2} |\tilde{p}((\kappa + i\eta)\beta)| \rightarrow 0$  by the Riemann-Lebesgue lemma. This, the continuity of  $|\tilde{p}((\kappa + i\eta)\beta)|$  with respect to  $\eta$ , and relation (27), imply the existence of  $\eta = \eta(\kappa)$ , such that equality (28) holds, and, consequently, inequality (26) holds. This  $\eta(\kappa)$  satisfies (29) because  $\mathcal{P}$  is bounded.

Lemma 4 is proved.  $\square$

To complete the proof of Theorem 1 one has to establish estimate (25). This estimate will be established if one proves the following relation:

$$\lim_{\kappa \rightarrow \infty} \nu(\kappa) := \lim_{\kappa \rightarrow \infty} \nu(\kappa, \eta(\kappa)) = 0, \quad (43)$$



where  $\eta(\kappa)$  satisfies (29) and

$$\nu(\kappa, \eta) = \sup_{\beta \in S^2} \int_{\mathbb{R}^3} |\tilde{\epsilon}((\kappa + i\eta)\beta - s)| ds. \quad (44)$$

Our argument is valid for  $\epsilon_1$ ,  $\epsilon_2$  and  $\epsilon_1\epsilon_2$ , so we will use the letter  $\epsilon$  and equation (13) for  $\tilde{\epsilon}$ .

Below we denote  $2k := \kappa + i\eta$  and we choose  $\eta = \eta(\kappa) = a^{-1} \ln \kappa + O(1)$  as  $\kappa \rightarrow \infty$ .

We prove that equation (12) can be solved by iterations if  $\text{Im}\eta \geq 0$  and  $|k + i\eta|$  is sufficiently large, because for such  $k + i\eta$  the operator  $T^2$  has small norm in  $C(B_a)$ , the space of functions, continuous in the ball  $B_a$ , with the sup-norm. Since equation (12) can be solved by iterations and the norm of  $T^2$  is small, the main term in the series, representing its solution, as  $|\kappa + i\eta| \rightarrow \infty$ ,  $\eta \geq 0$ , is the free term of the equation (12). The same is true for the Fourier transform of equation (12), i.e., for equation (13). Therefore the main term of the solution  $\tilde{\epsilon}$  to equation (13) as  $|\kappa + i\eta| \rightarrow \infty$ ,  $\eta \geq 0$ , is obtained by using the estimate of the free term of this equation. Thus, it is sufficient to check estimate (43) for the function  $\nu(\kappa, \eta(\kappa))$  using in place of  $\tilde{\epsilon}$  the function  $\tilde{q}(\xi)(\xi^2 - 2k\beta \cdot \xi)^{-1}$ , with  $2k$  replaced by  $\kappa + i\eta$  and  $\eta = a^{-1} \ln \kappa + O(1)$  as  $\kappa \rightarrow \infty$ .

For the above claim that equation (12) has the operator

$$T\epsilon = \int_{B_a} G(x - y, k)q(y)\epsilon(y, \beta, k)dy,$$

with the norm  $\|T^2\|$  in the space  $C(B_a)$ , which tends to zero as  $|\kappa + i\eta| \rightarrow \infty$ ,  $\eta \geq 0$ , see Appendix.

Thus, let us estimate the modulus of the factor  $\nu(\kappa, \eta)$  in (24) with  $\eta = \eta(\kappa)$  as in (29). Using inequality (40), and denoting  $\xi = (\kappa + i\eta)\beta$ , where  $\beta \in S^2$  plays the role of  $\alpha$  in (13), one obtains:

$$\begin{aligned} I &:= \sup_{\beta \in S^2} \int_{\mathbb{R}^3} \frac{|\tilde{q}((\kappa + i\eta)\beta - s)| ds}{|[(\kappa + i\eta)\beta - s]^2 - (\kappa + i\eta)\beta \cdot ((\kappa + i\eta)\beta - s)|} \\ &\leq ce^{a|\eta|} \sup_{\beta \in S^2} \int_{\mathbb{R}^3} \frac{ds}{|s^2 - (\kappa + i\eta)\beta \cdot s| [1 + (\kappa\beta - s)^2 + \eta^2]^{\ell/2}} \\ &:= ce^{a|\eta|} J. \end{aligned} \quad (45)$$

Let us prove that

$$J = o\left(\frac{1}{\kappa}\right), \quad \kappa \rightarrow \infty.$$

If this estimate is proved and  $\eta = a^{-1} \ln \kappa + O(1)$ , then  $I = o(1)$  as  $\kappa \rightarrow \infty$ , therefore relation (43) follows, and Theorem 1 is proved.

Let us write the integral  $J$  in the spherical coordinates with  $x_3$ -axis directed along vector  $\beta$ . We have

$$|s| = r, \quad \beta \cdot s = r \cos \theta := rt, \quad -1 \leq t \leq 1.$$

Denote

$$\gamma := \kappa^2 + \eta^2.$$

Then

$$\begin{aligned} J &\leq 2\pi \int_0^\infty dr r \int_{-1}^1 \frac{dt}{[(r - \kappa t)^2 + \eta^2 t^2]^{1/2} (1 + \gamma + r^2 - 2r\kappa t)^{\ell/2}} \\ &:= 2\pi \int_0^\infty dr r B(r), \end{aligned} \quad (46)$$

where

$$B := B(r) = B(r, \kappa, \eta) := \int_{-1}^1 \frac{dt}{[(r - \kappa t)^2 + \eta^2 t^2]^{1/2} (1 + \gamma + r^2 - 2r\kappa t)^{\ell/2}}.$$

Estimate of  $J$  we start with the observation

$$\tau := \min_{t \in [-1, 1]} [(r - \kappa t)^2 + \eta^2 t^2] = \min\{r^2 \eta^2 / \gamma, (r - \kappa)^2 + \eta^2\}.$$

Let  $\tau = r^2 \eta^2 / \gamma$ , which is always the case if  $r$  is sufficiently small. In the case when  $\tau = (r - \kappa)^2 + \eta^2$  the proof is considerably simpler and is left for the reader. If  $\tau = r^2 \eta^2 / \gamma$ , then

$$J \leq 2\pi \gamma^{1/2} \eta^{-1} \int_0^\infty dr \int_{-1}^1 dt [1 + \gamma + r^2 - 2\kappa r t]^{-\ell/2}.$$

Integrating over  $t$  yields

$$J \leq 2\pi \gamma^{1/2} \eta^{-1} [(\ell - 2)\kappa]^{-1} \mathcal{J},$$

where

$$\mathcal{J} := \int_0^\infty dr r^{-1} [(1 + \gamma + r^2 - 2\kappa r)^{-b} - (1 + \gamma + r^2 + 2\kappa r)^{-b}],$$

and  $b := \ell/2 - 1$ .

Since  $\eta = O(\ln \kappa)$ , one has  $\frac{\eta}{\kappa} = o(1)$  as  $\kappa \rightarrow \infty$ . Therefore,

$$\gamma^{1/2} \eta^{-1} \kappa^{-1} = O(\eta^{-1}) \quad \text{as } \kappa \rightarrow \infty.$$

Since  $\ell > 3$ , one has  $b > \frac{1}{2}$ , and, as we prove below,

$$\mathcal{J} = o\left(\frac{1}{\kappa}\right) \quad \text{as } \kappa \rightarrow \infty. \quad (47)$$

This relation implies the desired inequality:

$$J \leq o\left(\frac{1}{\kappa}\right) \quad \text{as } \kappa \rightarrow \infty. \quad (48)$$

Let us derive relation (47). One has

$$\mathcal{J} = \int_0^1 + \int_1^\infty := J_1 + J_2,$$

$$J_1 \leq \int_0^1 dr r^{-1} \frac{(w^2 + 2r\kappa + r^2)^b - (w^2 - 2r\kappa + r^2)^b}{(w^2 + 2r\kappa + r^2)^b (w^2 - 2r\kappa + r^2)^b},$$

where

$$w^2 := 1 + \gamma = 1 + \eta^2 + \kappa^2.$$

Furthermore,

$$(w^2 + 2r\kappa + r^2)^b - (w^2 - 2r\kappa + r^2)^b \leq \frac{4br\kappa}{(w^2 - 2r\kappa + r^2)^{1-b}}.$$

Thus,

$$J_1 \leq 4b\kappa \int_0^1 dr \frac{1}{(w^2 + 2r\kappa + r^2)^b (w^2 - 2r\kappa + r^2)}.$$

This implies the following estimate

$$J_1 \leq O(\kappa/w^{2+2b}) \leq O(\kappa^{-(1+2b)}),$$

because  $w = \kappa[1 + o(1)]$  as  $\kappa \rightarrow \infty$ . Furthermore,

$$J_2 \leq \int_1^\infty dr r^{-1} [(1 + \eta^2 + (r - \kappa)^2)^{-b} - (1 + \eta^2 + (r + \kappa)^2)^{-b}] := J_{21} - J_{22}.$$

One has  $J_{22} \leq J_{21}$ .

Let us estimate  $J_{21}$ . One obtains

$$J_{21} = \int_1^{\kappa/2} + \int_{\kappa/2}^\infty := j_1 + j_2,$$

and

$$j_1 \leq \frac{1}{[W^2 + \frac{\kappa^2}{4}]^b} \ln \kappa = o\left(\frac{1}{\kappa}\right), \quad W^2 := 1 + \eta^2, \quad b > \frac{1}{2}.$$

Furthermore

$$j_2 \leq \frac{2}{\kappa} \int_{\kappa/2}^\infty \frac{dr}{[W^2 + (r - \kappa)^2]^b} \leq \frac{2}{\kappa} \int_{-\infty}^\infty \frac{dy}{[W^2 + y^2]^b} = o\left(\frac{1}{\kappa}\right).$$

Thus, if  $b > \frac{1}{2}$ , then  $J_2 = o\left(\frac{1}{\kappa}\right)$  and  $\mathcal{J} = J_1 + J_2 = o\left(\frac{1}{\kappa}\right)$ . Thus, relation (47) is proved.

Relation (47) yields the desired estimate

$$J = o\left(\frac{1}{\kappa}\right).$$

Thus, both estimates (47) and (48) are proved.

Note that the desired relation  $\mathcal{J} = o\left(\frac{1}{\kappa}\right)$  could have been obtained even by replacing  $W^2$  by the smaller quantity 1 in the above argument.

Estimate (45) implies

$$I \leq ce^{a|\eta|} o\left(\frac{1}{\sqrt{\kappa^2 + \eta^2}}\right), \quad \kappa \rightarrow \infty, \quad \eta = a^{-1} \ln \kappa + O(1). \quad (49)$$

The quantity  $\eta = \eta(k) = a^{-1} \ln \kappa + O(1)$  was chosen so that if  $\kappa \rightarrow \infty$ , then the quantity  $\frac{e^{|\eta|a}}{\sqrt{\kappa^2 + \eta^2}}$  remains bounded as  $\kappa \rightarrow \infty$ . Therefore estimate (49) implies

$$\lim_{\kappa \rightarrow \infty, \eta = a^{-1} \ln \kappa + O(1)} I = 0. \quad (50)$$

Consequently, estimate (43) holds.

Theorem 1 is proved.  $\square$

## APPENDIX

### 1. Estimate of the norm of the operator $T^2$ .

Let

$$Tf := \int_{B_a} G(x - y, \kappa + i\eta) q(y) f(y) dy. \quad (51)$$

Assume  $q \in H_0^\ell(B_a)$ ,  $\ell > 2$ ,  $f \in C(B_a)$ . Our goal is to prove that equation (12) can be solved by iterations for all sufficiently large  $\kappa$ .

Consider  $T$  as an operator in  $C(B_a)$ . One has:

$$\begin{aligned} T^2 f &= \int_{B_a} dz G(x - z, \kappa + i\eta) q(z) \int_{B_a} G(z - y, \kappa + i\eta) q(y) f(y) dy \\ &= \int_{B_a} dy f(y) q(y) \int_{B_a} dz q(z) G(x - z, \kappa + i\eta) G(z - y, \kappa + i\eta). \end{aligned} \quad (52)$$

Let us estimate the integral

$$\begin{aligned} I(x, y) &:= \int_{B_a} G(x - z, \kappa + i\eta) G(z - y, \kappa + i\eta) q(z) dz \\ &= \int_{B_a} \frac{e^{i(\kappa+i\eta)[|x-z|-\beta \cdot (x-z)+|z-y|-\beta \cdot (z-y)]}}{16\pi^2 |x-z||z-y|} q(z) dz \\ &= \frac{1}{16\pi^2} \int_{B_a} \frac{e^{i(\kappa+i\eta)[|x-z|+|z-y|-\beta \cdot (x-y)]}}{|x-z||z-y|} q(z) dz \\ &:= \frac{e^{-i(\kappa+i\eta)\beta \cdot (x-y)}}{16\pi^2} I_1(x, y). \end{aligned} \quad (53)$$

Let us use the following coordinates (see [11], p.391):

$$z_1 = \ell st + \frac{x_1 + y_1}{2}, \quad z_2 = \ell \sqrt{(s^2 - 1)(1 - t^2)} \cos \psi + \frac{x_2 + y_2}{2}, \quad (54)$$

$$z_3 = \ell \sqrt{(s^2 - 1)(1 - t^2)} \sin \psi + \frac{x_3 + y_3}{2}. \quad (55)$$

The Jacobian  $J$  of the transformation  $(z_1, z_2, z_3) \rightarrow (\ell, t, \psi)$  is

$$J = \ell^3 (s^2 - t^2), \quad (56)$$

where

$$\ell = \frac{|x - y|}{2}, \quad |x - z| + |z - y| = 2\ell s, \quad |x - z| - |z - y| = 2\ell t, \quad (57)$$

$$|x - z||z - y| = 4\ell^2 (s^2 - t^2), \quad 0 \leq \psi < 2\pi, \quad t \in [-1, 1], \quad s \in [1, \infty). \quad (58)$$

One has

$$I_1 = \ell \int_a^\infty e^{2i(\kappa + i\eta)\ell s} Q(s) ds, \quad (59)$$

where

$$Q(s) := Q(s, \ell, \frac{x + y}{2}) = \int_0^{2\pi} d\psi \int_{-1}^1 dt q(z(s, t, \psi; \ell, \frac{x + y}{2})), \quad (60)$$

and the function  $Q(s) \in H_0^2(\mathbb{R}^3)$  for any fixed  $x, y$ . Therefore, an integration by parts in (59) yields the following estimate:

$$|I_1| = O\left(\frac{1}{|\kappa + i\eta|}\right), \quad |\kappa + i\eta| \rightarrow \infty. \quad (61)$$

From (52), (53) and (61) one gets:

$$\|T^2\| = O\left(\frac{1}{\sqrt{\gamma}}\right), \quad \gamma := \kappa^2 + \eta^2 \rightarrow \infty. \quad (62)$$

Therefore, integral equation (12), with  $k$  replaced by  $\frac{\kappa + i\eta}{2}$ , can be solved by iterations if  $\gamma$  is sufficiently large and  $\eta \geq 0$ . Consequently, integral equation (13) can be solved by iterations. Thus, estimate (43) holds if such an estimate holds for the free term in equation (13), that is, for the function  $\frac{\bar{q}}{\xi^2 - (\kappa + i\eta)\beta \cdot \xi}$ , namely, if estimate (50) holds.

## 2. Proof of Lemma 1.

Let  $L_j G_j := [\nabla^2 + k^2 - q_j(x)]G_j(x, y, k) = -\delta(x - y)$  in  $\mathbb{R}^3$ ,  $j = 1, 2$ . Applying Green's formula one gets

$$G_1(x, y, k) - G_2(x, y, k) = \int_{B_a} [q_2(z) - q_1(z)]G_1(x, z, k)G_2(z, y, k) dz. \quad (63)$$

In [11], p. 46, the following formula is proved:

$$G_j(x, y, k) = \frac{e^{ik|y|}}{4\pi|y|} u_j(x, \alpha, k) + o\left(\frac{1}{|y|}\right), \quad |y| \rightarrow \infty, \alpha := -\frac{y}{|y|}, \quad (64)$$

where  $u_j(x, \alpha, k)$  is the scattering solution,  $j = 1, 2$ . Applying formula (64) to (63), one obtains

$$u_1(x, \alpha, k) - u_2(x, \alpha, k) = \int_{B_a} [q_2(z) - q_1(z)]G_1(x, z, k)u_2(z, \alpha, k) dz \quad (65)$$

using the definition (2) of the scattering amplitude  $A(\beta, \alpha, k)$ , one derives from (65) the relation

$$4\pi[A_1(\beta, \alpha, k) - A_2(\beta, \alpha, k)] = \int_{B_a} [q_2(z) - q_1(z)]u_1(z, -\beta, k)u_2(z, \alpha, k)dz. \quad (66)$$

This formula is equivalent to (14) because of the well-known reciprocity relation  $A(\beta, \alpha, k) = A(-\alpha, -\beta, k)$ .

Lemma 1 is proved. □

## References

- [1] B. Levin, *Distribution of zeros of entire functions*, AMS, Providence, RI, 1980.
- [2] G. Polya, G. Szegő, *Problems and theorems in analysis*, Springer Verlag, Berlin, 1983, Vol.1, problem III.6.5.322.
- [3] A.G. Ramm, *Uniqueness theorem for inverse scattering with non-overdetermined data*. J. Phys. A, 43 (2010), 112001.
- [4] A.G. Ramm, *On the analytic continuation of the solution of the Schrödinger equation in the spectral parameter and the behavior of the solution to the nonstationary problem as  $t \rightarrow \infty$* . Uspechi Mat. Nauk, 19 (1964), 192 – 194 (in Russian).
- [5] A.G. Ramm, *Some theorems on analytic continuation of the Schrödinger operator resolvent kernel in the spectral parameter*. Izvestiya Acad. Nauk Armyan. SSR, Mathematics, 3 (1968), 443 – 464 (in Russian).
- [6] A.G. Ramm, *Recovery of the potential from fixed energy scattering data*. Inverse Problems, 4 (1988), 877 – 886; 5 (1989) 255.
- [7] A.G. Ramm, *Stability estimates in inverse scattering*. Acta Appl. Math., 28, no. 1 (1992), 1 – 42.
- [8] A.G. Ramm, *Stability of the solution to inverse obstacle scattering problem*. J.Inverse and Ill-Posed Problems, 2, no. 3 (1994), 269 – 275.
- [9] A.G. Ramm, *Stability of solutions to inverse scattering problems with fixed-energy data*. Milan Journ of Math., 70, (2002), 97 – 161.
- [10] A.G. Ramm, *Inverse problems*, Springer, New York, 2005.
- [11] A.G. Ramm, *Scattering by obstacles*, D.Reidel, Dordrecht, 1986.
- [12] A.G. Ramm, A.I. Katsevich, *The Radon transform and local tomography*, CRC Press, Boca Raton, 1996.

Alexander Ramm  
Department of Mathematics  
Kansas State University  
Manhattan, KS 66506-2602, USA  
E-mail: ramm@math.ksu.edu

Received: 29.04.2010